CYP3A and CYP2CI9 activity in urine in relation to CYP3A4, CYP3A5, and CYP2CI9 polymorphisms in Russian peptic ulcer patients taking omeprazole

Natalia P Denisenko¹⁻³ Dmitriy A Sychev² Zhanna M Sizova³ Valeriy V Smirnov^{4,5} Kristina A Ryzhikova¹ Zhannet A Sozaeva¹ Elena A Grishina¹

¹Research Center, Russian Medical Academy of Continuous Professional Education, Ministry of Healthcare, Moscow, Russia; ²Department of Clinical Pharmacology and Therapy, Russian Medical Academy of Continuous Professional Education, Ministry of Healthcare, Moscow, Russia; 3Department of Social Expertise, Urgent and Outpatient Therapy, First Moscow State Medical University (Sechenov University), Ministry of Healthcare, Moscow, Russia; ⁴Department of Pharmaceutical and Toxicological Chemistry, First Moscow State Medical University (Sechenov University), Ministry of Healthcare, Moscow, Russia; 5Laboratory of Clinical Pharmacology, National Research Centre - Institute of Immunology, Federal Medical Biological Agency, Moscow, Russia

Background: Proton pump inhibitors (PPIs) are metabolized by cytochrome P450. CYP2C19 is the main isoenzyme for the majority of PPI, whereas CYP3A family is a secondary enzyme for PPI biotransformation.

Purpose: The aim of the study was to find if CYP3A4*22, CYP3A5*3, CYP2C19*2, CYP2C19*3, and CYP2C19*17 genotypes are connected with CYP3A and CYP2C19 activities in Russian peptic ulcer patients taking omeprazole.

Patients and methods: Forty-eight gastric or duodenal ulcer patients (15 men, 33 women; mean age 55.0±15.3 years, age range 18-91 years) from Moscow region of Russia were enrolled. Peripheral venous blood was collected for DNA extraction, and real-time polymerase chain reaction was performed for CYP3A5*3^{A6986G} (rs776746), CYP3A4*22 C>T in intron 6 (rs35599367), CYP2C19*2^{G681A} (rs4244285), CYP2C19*3^{G636A} (rs4986893), and CYP2C19*17^{C-806T} (rs12248560) polymorphism analyses. Urine samples of patients were collected in the morning between 6 and 9 am before food or drug intake. Urine cortisol and 6β-hydroxycortisol concentrations (for CYP3A activity) and omeprazole and 5-hydroxyomeprazole concentrations (for CYP2C19 activity) were measured using high-performance liquid chromatography/mass spectroscopy.

Results: We found a connection between CYP2C19 genotypes and CYP3A activity. Median metabolic ratios 6β-hydroxycortisol/cortisol (25%-75% percentiles) were 2.84 (1.99-4.39) for CYP2C19 extensive metabolizers (EMs), 2.51 (1.86-4.73) for CYP2C19 ultra-rapid metabolizers (UMs), and 1.45 (1.12–2.16) for CYP2C19 intermediate metabolizers (IMs) + poor metabolizers (PMs). A statistically significant difference in CYP3A activity (Mann-Whitney test) was found between CYP2C19 EMs vs CYP2C19 IMs+PMs (p=0.006), between CYP2C19 UMs vs CYP2C19 IMs+PMs (p=0.018), and in multiple comparison Kruskal–Wallis test (p=0.014). Conclusion: In CYP2C19 IMs+PMs, CYP3A activity was significantly lower than in CYP2C19 EMs and UMs.

Keywords: pharmacogenetics, phenotyping, metabolomics, proton pump inhibitor

Introduction

Proton pump inhibitors (PPIs), which are widely used in peptic ulcer patients' treatment, are metabolized by cytochrome P450. CYP2C19 is the main isoenzyme for the majority of PPI, whereas CYP3A family is a secondary enzyme for PPI biotransformation. 1 CYP2C19 polymorphisms are studied in detail and their impact on PPI efficacy is established in guidelines of Dutch Pharmacogenetics Working Group of the Royal Dutch Pharmacists Association.²

Correspondence: Natalia P Denisenko 125993, Barrikadnaja Street, 2/1, Moscow, Russia Tel +7 495 945 8139 Email natalypilipenko3990@gmail.com CYP3A4 and CYP3A5 are the main significant isoenzymes of CYP3A family in adults. CYP3A4 metabolizes about 60% of all known drugs. Metabolism of PPI involves CYP3A enzymes, and CYP3A may become the main pathway especially in CYP2C19 poor metabolizers (PMs). CYP3A4 activity may be detected using 6 β -hydroxycortisol/cortisol ratios in urine. CYP3A4 genetic polymorphisms are being studied. CYP3A4*22, found in 2011, is associated with low CYP3A4 expression and low CYP3A4 activity. CYP3A4*22 allele influences efficacy of tacrolimus, statin, and cyclosporine therapy. β -11

It was shown that CYP3A5 and CYP3A4 have common substrates. 12-14 Studies showed that CYP3A5 genotypes reflect CYP3A4 activity. 12 So, CYP3A5 genetic polymorphisms may also have impact on PPI metabolism. 15 It was shown that CYP3A5*3 allele is associated with lack of activity of the enzyme compared with wild-type CYP3A5*1 allele. 3

CYP3A phenotype may be predicted on the basis of CYP3A4*22 and CYP3A5*3 carrier. Persons with CYP3A4*1/*1 genotype (no CYP3A4*22) together with CYP3A5*1 carrier were classified as CYP3A EMs. CYP3A IMs were CYP3A4*22 together with CYP3A5*1 carrier or CYP3A4*1/*1 with CYP3A5*3/*3 genotypes. CYP3A PMs were CYP3A4*22 and CYP3A5*3/*3 carriers. ¹⁶

The aim of the study was to find if CYP3A4*22, CYP3A5*3, CYP2C19*2, CYP2C19*3, and CYP2C19*17 genotypes are connected with CYP3A and CYP2C19 activities in Russian peptic ulcer patients.

Patients and methods Study design and patients

Forty-eight gastric or duodenal ulcer patients (15 men, 33 women; mean age 55.0±15.3 years, age range 18–91 years) from Moscow region of Russia were enrolled in the study after providing written informed consent to participate in the study. This study is an extension of previously published study.¹⁷

Peptic ulcer diagnosis was based on endoscopy and histology. There were 18 gastric ulcer patients, 25 duodenal ulcer patients, and 5 patients had both gastric and duodenal ulcers. Patients underwent gastroscopy and were prescribed omeprazole 20 mg BID for at least 7 days intake. Exclusion criteria were severe comorbidities (cardiovascular, cerebral, renal, hepatic dysfunction) and CYP3A inhibitor clarithromycin

intake. The study was approved by local Ethics Committee of First Moscow State Medical University (Sechenov University, Moscow, Russia).

Statins (atorvastatin or simvastatin), which refer to weak CYP3A inducers, were prescribed to 3 patients.

Genotyping

Peripheral venous blood (6 mL) of patients was collected in K2-EDTA tubes for DNA extraction from leukocytes. Blood samples were stored in the deep freeze at –70°C until analysis. DNA was isolated using a commercially available kit "S-sorb" (produced by "Syntol," Moscow, Russia) with silica-based sorbent. *CYP3A5*3* ^{A6986G} (rs776746) and *CYP3A4*22* C>T in intron 6 (rs35599367) polymorphisms were analyzed using real-time polymerase chain reaction (PCR). PCR was performed using a commercially available kit (produced by "Syntol") in DNA amplifier CFX96 Touch (Bio-Rad Laboratories, Hercules, CA, USA) for *CYP3A5*3* and TaqMan genotyping assay (produced by Applied Biosystems Inc., Foster City, CA, USA) for *CYP3A4*22*.

The program consisted of initial denaturation step at 95°C for 3 minutes followed by 40 cycles of 15 seconds' denaturation at 95°C, then annealing at 63°C for 40 seconds. Genotyping was repeated for 20% of patients' samples as a control for correct sample handling. Repeated genotyping revealed identical results.

CYP2C19*2, CYP2C19*3, and CYP2C19*17 genotyping have been described previously.¹⁷

Assay

Urine samples were stored in deepfreeze at -70° C until analysis. About 2 mL of urine was extracted with 4 mL of ethyl acetate/isopranolol (85:15). After orbital mixing for 10 minutes and centrifuging at 3,000×g for 5 minutes, the upper organic layer was separated and transferred to glass tube. About 2 mL 1 M sodium hydroxide solution was added to organic layer, then followed orbital mixing for 10 minutes. and centrifuging at 3,000×g for 5 minutes to separate organic layer and evaporation to dryness under gentle stream of air was finished dissolving dry residue in 1 mL of ethanol.

Phenotyping

Endogenous cortisol transforms to 6 β -hydroxycortisol selectively by CYP3A isoenzyme; thus, the metabolic ratio of 6 β -hydroxycortisol/cortisol in urine is suggested to reflect the activity of CYP3A. The higher the urine metabolic ratio

of 6β -hydroxycortisol/cortisol, the higher the activity of CYP3A. As cortisol concentration is the highest in the morning, urine samples of patients were collected between 6 and 9 am before food or drug intake.

Cortisol and 6\beta-hydroxycortisol concentrations in urine were measured using Agilent 1290 Infinity (Agilent Technologies, Santa Clara, CA, USA) high-performance liquid chromatography with mass spectrometry. The isolation of drug and its metabolite was performed on Waters Symmetry C18 Column (150×4.6 mm; 5.0 µm, Waters Corporation, Milford, MA, USA). The column temperature was maintained at 35°C. UV detector wavelength was set at 246 nm. The mobile phase contained of 55% water formic acid solution (1 L of water:1 mL of formic acid) and 45% acetonitrile. The flow rate was 0.5 mL/min. Volumes of 10 μL were injected. The mass spectrometer was operated using the following conditions: positive polarity, MM-ES+APCI ionization.

CYP2C19 phenotyping using omeprazole and 5-hydroxyomeprazole concentrations in urine was described previously.¹⁷

Statistical analysis

The statistical analysis was performed using SPSS Statistics 22 (IBM Corporation, Armonk, NY, USA). A p-value < 0.05 was considered statistically significant. The normality of metabolic ratio distribution was analyzed by Kolmogorov-Smirnov test. Nonparametric statistics was used to compare metabolic ratio in different genotypes. The observed and expected genotype frequencies were calculated by using Hardy-Weinberg equilibrium. Fisher's exact test was used to evaluate deviation of genotype frequencies in the studied population from Hardy-Weinberg equilibrium.

Results CYP2C19, CYP3A4, and CYP3A5 genotyping results

Of 48 patients, there were 46 (95.8%) CC (CYP3A4*1/*1), 1 (2.1%) CT (CYP3A4*1/*22), and 1 (2.1%) TT (CYP3A4*22/*22) carriers of CYP3A4*22 (Table 1). Genotypes were not in accordance to Hardy-Weinberg equilibrium $(\chi^2=20.7, p=0.00)$. The CYP3A4*22 allele frequency was 3.1%.

Of 48 patients, there were 4(8.3%) GA (CYP3A5*1/*3) and 44 (91.7%) GG (CYP3A5*3/*3) carriers of CYP3A5*3 A6986G (Table 2). Genotypes were in accordance to Hardy-Weinberg equilibrium, p>0.05 ($\chi^2=0.09$, p=0.76). The CYP3A5*3 allele frequency was 95.8%.

Of 48 patients, there were 4 (8.3%) CYP3A extensive metabolizers (EMs), 42 (87.5%) CYP3A intermediate metabolizers (IMs), and 2 (4.2%) CYP3A PMs (Table 3).

Results of CYP2C19 genotyping for 59 patients were described previously in detail.¹⁷ Eleven patients from the previous study were excluded for the current analysis because of clarithromycin, a CYP3A inhibitor, intake. Results of CYP2C19 genotyping for 48 patients are described in Tables 2 and 3.

Table 3 CYP3A4 and CYP3A5 genotypes and predicted CYP3A phenotype in Russian peptic ulcer patients

СҮРЗА	CYP3A4*22	CYP3A5*3	Total,	
phenotype	genotype	genotype	patients	
EMs	СС	GA	4	
IMs	CT+TT	AA+GA	_	
	CC	GG	42	
PMs	CT+TT	GG	2	
Total			48	

Abbreviations: EMs. extensive metabolizers: IMs. intermediate metabolizers: PMs. poor metabolizers.

Table I CYP3A4 and CYP2C19 genotypes in Russian peptic ulcer patients

CYP3A4*22 C>T in intron 6	CYP2C19							Total,
	*1/*1	*1/*17	*1/*2	*2/*17	*2/*2	*3/*17	*17/*17	patients
СС	19	13	8	2	2	I	I	46
CT	_	1	_	_	_	_	_	1
TT	1	_	_	_	_	_	_	1
Total	20	14	8	2	2	1	1	48

Table 2 CYP3A5 and CYP2C19 genotypes in Russian peptic ulcer patients

CYP3A5 A6986G	CYP2C19							Total,
	*1/*1	*1/*17	*1/*2	*2/*17	*2/*2	*3/*17	*17/*17	patients
GG	19	13	7	2	I	1	I	44
GA	1	1	1	_	1	_	_	4
Total	20	14	8	2	2	1	1	48

CYP3A activity and its relationship to CYPs genotypes

Because distribution of metabolic ratio of 6β -hydroxycortisol/cortisol was not normal (Kolmogorov–Smirnov Test p=0.001) median and 25%–75% percentiles for data description and nonparametric statistics for samples' comparison were used.

CYP3A4*22

Median metabolic ratio of 6β-hydroxycortisol/cortisol for all genotypes was 2.30 (1.47–4.13). Median 6β-hydroxycortisol/cortisol ratio for CYP3A4*1/*1 was 2.27 (1.46–3.70), and for CYP3A4*1/*22+CYP3A4*22/*22, it was 8.65.

We found statistically significant difference in urine metabolic ratio of 6β-hydroxycortisol/cortisol between patients with *CYP3A4*1/*1* genotype vs *CYP3A4*1/*22* +*CYP3A4*22/*22* carriers (Mann–Whitney test, *p*=0.03). So, we found that CYP3A activity was significantly higher in *CYP3A4*22* carriers, which contadicts the data of previous studies.

CYP3A5*3

Median metabolic ratio of 6β -hydroxycortisol/cortisol for CYP3A5*1/*3 was 2.27 (1.45–4.20), and for CYP3A5*3/*3, it was 2.27 (1.40–4.34).

No statistically significant difference in CYP3A activity (Mann–Whitney test) was found between CYP3A5*1/*3 vs CYP3A5*3/*3 carriers (p=0.63).

CYP2C19*2, *3, and *17

Patients were referred to as *CYP2C19* EMs, PMs, IMs, or ultra-rapid metabolizers (UMs) according to the Dutch Pharmacogenetics Working Group Guideline of the Royal Dutch Pharmacists Association.² Median metabolic ratio of 6β -hydroxycortisol/cortisol for CYP2C19 EMs was 2.84 (1.99–4.39), UMs – 2.51 (1.86–4.73), and IMs+PMs – 1.45 (1.12–2.16).

Statistically significant difference in CYP3A activity (Mann–Whitney test) was found between EMs vs IMs+PMs (p=0.006), between UMs vs IMs+PMs (p=0.018), and in multiple comparison Kruskal–Wallis test (p=0.014) (Table 4).

Both CYP3A4*22 and CYP3A5*3

When comparing CYP3A activity between CYP3A EMs and a combined group of CYP3A IMs+PMs, the results correspond to the comparison of CYP3A5*3 genotypes with no statistically significant difference between groups (*p*=0.63).

CYP2C19 activity and its relationship to CYPs genotypes

CYP2C19*2, *3, and *17

The results have been previously published.¹⁷

CYP3A4*22

Median metabolic ratio of 5-hydroxyomeprazole/omeprazole in urine for all genotypes was 1.23 (0.80–1.99); for CYP3A4*1/*1, it was 1.23 (0.79–1.88), and for CYP3A4*1/*22+CYP3A4*22/*22, it was 1.62.

No statistically significant difference in CYP2C19 activity (Mann–Whitney test) was found between *CYP3A4*1/*1* vs *CYP3A4*1/*22+CYP3A4*22/*22* carriers (*p*=0.76).

CYP3A5*3

Median metabolic ratio of 5-hydroxyomeprazole/omeprazole in urine for *CYP3A5*1/*3* was 2.40 (0.76–2.98) and for *CYP3A5*3/*3* was 1.17 (0.80–1.79).

No statistically significant difference in CYP2C19 activity (Mann–Whitney test) was found between CYP3A5*1/*3 vs CYP3A5*3/*3 carriers (p=0.22).

Both CYP3A4*22 and CYP3A5*3

When comparing CYP2C19 activity between CYP3A EMs and a combined group of CYP3A IMs+PMs, the results

Table 4 CYP2C19 genotypes and CYP3A activity and in Russian peptic ulcers patients

Enzyme activity	CYP2C19 genotype									
	EMs		UMs		IMs+PMs		Total			
	Median	Percentiles 25%–75%	Median	Percentiles 25%–75%	Median	Percentiles 25%–75%	Median	Percentiles 25%–75%		
CYP3A activity	2.84	1.99-4.39	2.51	1.86–4.73	1.45	1.12–2.16	2.30	1.47–4.13		
	EMs vs UMs		EMs vs IMs+PMs		UMs vs IMs+PMs		EMs vs UMs vs IMs+PMs			
CYP3A activity	p=0.790		p=0.006		p=0.018		p=0.014			

Note: p<0.05 for bold text.

Abbreviations: EMs, extensive metabolizers; IMs, intermediate metabolizers; PMs, poor metabolizers; UMs, ultra-rapid metabolizers.

correspond to comparison of CYP3A5*3 genotypes with no statistically significant difference between groups (p=0.22).

Discussion

CYP3A5*3 allele is frequently seen among Caucasians (frequency 88%–97%), whereas in Hispanic and Asian population, it is more rare (66%–75%). As for African Americans and Africans, the frequency of CYP3A5*3 allele is about 37% or less. ¹⁸ Studies found 96.6%–97.7% CYP3A5*3 allele frequency in Russians. ^{19,20} We found 95.8% CYP3A5*3 allele frequency in our study of Russian peptic ulcer patients, which corresponds to data ^{19,20} described for Caucasians, and particularly Russians.

CYP3A4*22 allele is less frequent and is found in 5%–7% of Caucasians. ¹⁶ We found 3.1% CYP3A4*22 allele frequency in our study to be lower than data described for Caucasians, and CYP3A4*22 genotypes in our study were not in accordance with Hardy–Weinberg equilibrium.

CYP3A4 and CYP3A5 have overlapping substrate specificity, so it is important to genotype both CYP3A4 and CYP3A5 as well as to study CYP3A activity.

We did not find connections between CYP3A activity and CYP3A4 and CYP3A5 genotype, except the data that CYP3A activity was significantly higher in *CYP3A4*22* carriers. CYP3A activity was studied in some previous works, and no evidence was found for the fact that large interindividual differences in its activity were based on genotype. ^{21–24} In another study, midazolam and erythromycin were used for CYP3A phenotyping, and there were connections between CYP3A activity and CYP3A4*22 genotype and partly (only for midazolam) for CYP3A5*3 genotype. ¹⁶

We did not find connections between CYP2C19 activity and CYP3A4 and CYP3A5 genotypes, but found a connection between CYP2C19 genotype and CYP2C19 activity in a previously published study.¹⁷

We found a relationship between CYP2C19 genotype and CYP3A activity: in CYP2C19 IMs+PMs, CYP3A activity was significantly lower than in CYP2C19 EMs (p=0.001), and it also was significantly lower than in CYP2C19 UMs (p=0.011). The findings contradict our hypothesis that there is an increased CYP3A activity in CYP2C19 IMs and PMs compared to CYP2C19 EMs and UMs.

One of the limitations of our study was the following: we estimated CYP3A activity using 6β -hydroxycortisol/cortisol ratio in urine, not omeprazole sulfoxidation index (omeprazole sulfone/omeprazole in plasma), so it was not possible to identify the potential activation of the secondary metabolic route of conversion of omeprazole to omeprazole

sulfone in CYP2C19 IMs and PMs. Other limitations of the study were that only one single nucleotide polymorphism genotyping was done for CYP3A4 and also for CYP3A5, the low sample size, and the fact that CYP3A4*22 genotypes are not in Hardy–Weinberg equilibrium.

Conclusion

We found a connection between *CYP2C19* genotypes and CYP3A activity, measured using 6β-hydroxycortisol/cortisol ratio in urine in Russian peptic ulcer patients. In *CYP2C19* IMs+PMs, CYP3A activity was significantly lower than in *CYP2C19* EMs (*p*=0.006). Also, *CYP2C19* IMs+PMs had significantly lower CYP3A activity compared with *CYP2C19* UMs (*p*=0.018). There were no connections between CYP3A activity and *CYP3A5*3*. *CYP3A4*22* carriers had significantly higher CYP3A activity. No connections were found between CYP2C19 activity, measured using 5-hydroxyomeprazole/omeprazole ratio in urine and *CYP3A5*3* and *CYP3A4*22* genotypes.

Acknowledgment

This work is supported by Russian Scientific Foundation (project number 16-15-00227: "Fundamental research and exploratory research in priority areas of research") for CYP2C19*2 and CYP2C19*3 genotyping reagents and consumables and CYP2C19 and CYP3A phenotyping reagents and consumables and University Grant of First Moscow State Medical University (Sechenov University) ("Prognosing the efficacy of peptic ulcer therapy on the basis of pharmacogenomics, pharmacometabolomic and pharmacoproteomic technologies of personalized medicine") for CYP2C19*17, CYP3A4*22, and CYP3A5*5 genotyping reagents and consumables.

Disclosure

The authors report no conflicts of interest in this work.

References

- Li XQ, Weidolf L, Simonsson R, Andersson TB. Enantiomer/enantiomer interactions between the S- and R- isomers of omeprazole in human cytochrome P450 enzymes: major role of CYP2C19 and CYP3A4. *J Pharmacol Exp Ther*. 2005;315(2):777–787.
- Swen JJ, Nijenhuis M, de Boer A, et al. Pharmacogenetics: from bench to byte-an update of guidelines. Clin Pharmacol Ther. 2011;89(5):662-673.
- Kuehl P, Zhang J, Lin Y, et al. Sequence diversity in CYP3A promoters and characterization of the genetic basis of polymorphic CYP3A5 expression. *Nat Genet*. 2001;27(4):383–391.
- 4. Daly AK. Significance of the minor cytochrome P450 3A isoforms. *Clin Pharmacokinet*. 2006;45(1):13–31.
- Martinez-Jimenez CP, Jover R, Donato MT, Castell JV, Gomez-Lechon MJ. Transcriptional regulation and expression of CYP3A4 in hepatocytes. *Curr Drug Metab*. 2007;8(2):185–194.

- Shin KH, Ahn LY, Choi MH. Urinary 6β-hydroxycortisol/cortisol ratio most highly correlates with midazolam clearance under hepatic CYP3A inhibition and induction in females: a pharmacometabolomics approach. AAPS J. 2016;18(5):1254–1261.
- 8. Wang D, Guo Y, Wrighton SA, Cooke GE, Sadee W. Intronic polymorphism in CYP3A4 affects hepatic expression and response to statin drugs. *Pharmacogenomics J.* 2011;11(4):274–286.
- Elens L, Becker ML, Haufroid V, et al. Novel CYP3A4 intron 6 single nucleotide polymorphism is associated with simvastatin-mediated cholesterol reduction in the Rotterdam Study. *Pharmacogenet Genomics*. 2011;21(12):861–866.
- Elens L, Bouamar R, Hesselink DA, et al. A new functional CYP3A4 intron 6 polymorphism significantly affects tacrolimus pharmacokinetics in kidney transplant recipients. *Clin Chem.* 2011;57(11): 1574–1583.
- Passey C, Birnbaum AK, Brundage RC, et al. Validation of tacrolimus equation to predict troughs using genetic and clinical factors. *Pharma-cogenomics*. 2012;13(10):1141–1147.
- Lee SJ, Goldstein JA. Functionally defective or altered CYP3A4 and CYP3A5 single nucleotide polymorphisms and their detection with genotyping tests. *Pharmacogenomics*. 2005;6:357–371.
- Koch I, Weil R, Wolbold R, et al. Interindividual variability and tissue specificity in the expression of cytochrome P450 3A mRNA. *Drug Metab Dispos*. 2002;30(10):1108–1114.
- 14. Elens L, Bouamar R, Hesselink DA, Haufroid V, van Gelder T, van Schaik RH. The new CYP3A4 intron 6 C>T polymorphism (CYP3A4*22) is associated with an increased risk of delayed graft function and worse renal function in cyclosporine-treated kidney transplant patients. *Pharmacogenet Genomics*. 2012;22(5):373–380.

- Li Y, Zhang W, Guo D, Zhou G, Zhou H, Xiao Z. Pharmacokinetics of the new proton pump inhibitor ilaprazole in Chinese healthy subjects in relation to CYP3A5 and CYP2C19 genotypes. *Clin Chim Acta*. 2008;391(1–2):60–67.
- Elens L, van Gelder T, Hesselink DA, Haufroid V, van Schaik RH. CYP3A4*22: promising newly identified CYP3A4 variant allele for personalizing pharmacotherapy. *Pharmacogenomics*. 2013;14(1):47–62.
- Denisenko NP, Sychev DA, Sizova ZM, et al. Urine metabolic ratio of omeprazole in relation to CYP2C19 polymorphisms in Russian peptic ulcer patients. *Pharmgenomics Pers Med.* 2017;10:253–259.
- Zanger UM, Schwab M. Cytochrome P450 enzymes in drug metabolism: regulation of gene expression, enzyme activities, and impact of genetic variation. *Pharmacol Ther*. 2013;138(1):103–141.
- Mustafina OE, Tuktarova IA, Karimov DD, Somova Rs, Nasibullin TR. [CYP2D6, CYP3A5, and CYP3A4 gene polymorphisms in Russian, Tatar, and Bashkir populations]. *Genetika*. 2015;51(1):109–119. Russian.
- Zastrozhin MS, Grishina EA, Ryzhikova KA, et al. The influence of CYP3A5 polymorphisms on haloperidol treatment in patients with alcohol addiction. *Pharmgenomics Pers Med*. 2017;11:1–5.
- Lamba JK, Lin YS, Schuetz EG, Thummel KE. Genetic contribution to variable human CYP3A-mediated metabolism. *Adv Drug Deliv Rev.* 2002;54(10):1271–1294.
- Dai D, Tang J, Rose R, et al. Identification of variants of CYP3A4 and characterization of their abilities to metabolize testosterone and chlorpyrifos. *J Pharmacol Exp Ther*. 2001;299(3):825–831.
- Eiselt R, Domanski TL, Zibat A, et al. Identification and functional characterization of eight CYP3A4 protein variants. *Pharmacogenetics*. 2001;11(5):447–458.
- Ozdemir V, Kalow W, Tang BK, et al. Evaluation of the genetic component of variability in CYP3A4 activity: a repeated drug administration method. *Pharmacogenetics*. 2000;10(5):373–388.

Pharmacogenomics and Personalized Medicine

Publish your work in this journal

Pharmacogenomics and Personalized Medicine is an international, peerreviewed, open access journal characterizing the influence of genotype on pharmacology leading to the development of personalized treatment programs and individualized drug selection for improved safety, efficacy and sustainability. This journal is indexed on the American Chemical Society's Chemical Abstracts Service (CAS). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www.dovepress.com/testimonials.php to read real quotes from published authors.

Submit your manuscript here: https://www.dovepress.com/pharmacogenomics-and-personalized-medicine-journal

