

CYP2C19*17 May Increase the Risk of Death Among Patients with an Acute Coronary Syndrome and Non-Valvular Atrial Fibrillation Who Receive Clopidogrel and Rivaroxaban

This article was published in the following Dove Press journal:
Pharmacogenomics and Personalized Medicine

DA Sychev¹
OA Baturina¹ ²
KB Mirzaev¹
E Rytkin¹ ¹
DV Ivashchenko¹ ²
DA Andreev¹ ²
KA Ryzhikova¹ ¹
EA Grishina¹
PO Bochkov¹ ¹
RV Shevchenko¹ ¹

¹Russian Medical Academy of Continuous Professional Education, Ministry of Health of the Russian Federation, Moscow, Russian Federation; ²Sechenov University, Moscow, Russian Federation

Introduction: The aim of this study is to assess the influence of gene *CYP2C19*, *CYP3A4*, *CYP3A5* and *ABCB1* polymorphisms on clopidogrel antiplatelet activity, rivaroxaban concentration equilibrium, and clinical outcomes among patients with acute coronary syndrome and non-valvular atrial fibrillation.

Methods: In the multicenter prospective registry study of the efficacy and safety of a combined antithrombotic therapy 103 patients with non-valvular atrial fibrillation both undergoing or not a percutaneous coronary intervention were enrolled. The trial assessed the primary outcomes (major bleeding, in-hospital death, cardiovascular death, stroke/transient ischaemic attack, death/renal insufficiency) and secondary outcomes (platelet reactivity units (PRU), rivaroxaban concentration).

Results: For none of the clinical outcomes when combined with other covariates, the carriership of polymorphisms *CYP3A5*3 rs776746*, *CYP2C19*2 rs4244285*, **17 rs12248560*, *ABCB1 3435 C>T*, *ABCB1 rs4148738* was significant. None of the markers under study (*CYP3A5*3 rs776746*, *CYP2C19*2 rs4244285*, **17 rs12248560*, *ABCB1 3435 C>T*, *ABCB1 rs4148738*) has proven to affect rivaroxaban equilibrium concentration in blood plasma among patients with atrial fibrillation and acute coronary syndrome.

Conclusion: In situations of double or triple antithrombotic rivaroxaban and clopidogrel therapy among patients with atrial fibrillation and acute coronary syndrome, the genetic factors associated with bleeding complications risk (*CYP2C19*17*) may prove to be clinically relevant.

Keywords: rivaroxaban, clopidogrel, polymorphism, atrial fibrillation, acute coronary syndrome

Plain Language Summary

- A multicenter prospective registry trial was performed to study the efficacy and safety of a combined antithrombotic therapy: 103 patients with non-valvular atrial fibrillation some undergoing a percutaneous coronary intervention were included.
- The trial assessed the primary outcomes (major bleeding, in-hospital death, cardiovascular death, stroke/transient ischaemic attack, death/renal insufficiency) and secondary outcomes (platelet reactivity units, rivaroxaban concentration).
- In situations of double or triple antithrombotic rivaroxaban and clopidogrel therapy among patients with AF and ACS, the genetic factors associated with bleeding complications risk (*CYP2C19*17*) may prove to be clinically relevant.

Correspondence: E Rytkin
Russian Medical Academy of Continuous Professional Education, Ministry of Health of the Russian Federation, Moscow, Russian Federation
Email erytkin@gmail.com

Introduction

All the patients with an acute coronary syndrome regardless of the treatment strategy (with or without a percutaneous coronary intervention) were assigned to a double antiplatelet therapy, which includes aspirin and a P2Y12-inhibitor, in order to prevent thrombotic complications: cardiovascular death, myocardial infarction, stent thrombosis.¹ Nowadays, 3 P2Y12-inhibitors are available per os: clopidogrel, ticagrelor, prasugrel. In some trials the variability in response to clopidogrel was observed, which is connected with numerous factors, CYP2C19-dependent active metabolite production among them.^{2,3} Though the presence of low function alleles *CYP2C19*2* (*rs4244285*, *c.681G>A*) and *CYP2C19*3* (*rs4986893*, *c.636G>A*) leads to an insufficient platelet activity inhibition and significantly increases the risk of thrombotic complications compared to non-carriers.^{2,4} *CYP2C19*17* carriership is connected to an increased risk of bleeding complications on clopidogrel.³ The number of patients who are resistant to clopidogrel may be up to 35% of the population, which depends on the method of testing and the borders of high residual platelet reactivity.⁵ A genotype-based antiplatelet therapy, as observed in a number of observational and randomized trials, is capable of overcoming high residual platelet reactivity^{6–9} and may lead towards a decrease in the number of adverse cardiovascular events.^{8–16} These results are proven by the results of a meta-analysis of 9000 patients on clopidogrel: the carriership of low function allelic variants increases the risk of major adverse cardiovascular events (MACE) 1.5-fold and the risk of stent thrombosis 2.8-fold.¹⁷ The negative results of later meta-analyses can be explained by the heterogeneity of the population and inclusion in the meta-analyses of patients with a stable coronary disease.¹⁸ Meanwhile, several large randomized clinical trials are in progress (Popular genetics (NCT01761786) – 2700 patients; TAILOR-PCI (NCT01742117) – 5270 patients) and the clinical application of genotype-based antiplatelet therapy remains controversial. Other P2Y12-inhibitors – ticagrelor and prasugrel are not susceptible to the influence of allelic variants of the CYP2C19 gene carriership^{19,20} and have appeared to be more potent in terms of decrease in thrombotic complications in comparison with clopidogrel among patients with acute coronary syndrome undergoing percutaneous coronary intervention.^{21,22} Nevertheless, the problem of the variability of the response to clopidogrel is still relevant as the therapy with ticagrelor and prasugrel is connected to the higher number of bleeding complications; it is more expensive,

which leads to lower compliance among patients and, finally, is not the treatment of choice when used as part of a combined antithrombotic treatment.^{23,24}

According to the current guidelines and expert opinions, patients with an acute coronary syndrome (ACS) with atrial fibrillation (AF) need to be treated with combined antithrombotic therapy: antiplatelet therapy plus oral anticoagulants.²⁴ Although the prescription of anticoagulant therapy is regulated by the guidelines for treatment of AF aimed to prevent cardioembolic complications, a combined antithrombotic therapy is connected to an increase in the bleeding events risk.^{24–27} Nevertheless, the results of previous randomized clinical trials PIONEER AF-PCI, RE-DUAL PCI^{28,29} which included patients with ACS and AF undergoing PCI and meta-analyses^{30,31} showed the superiority of double antithrombotic treatment (oral anticoagulant+ P2Y12 inhibitor) over triple antithrombotic therapy in terms of decrease in the number of bleeding complications without significant differences in the effectiveness. Apart from AF treatment, prescription of oral anticoagulants may be pathophysiologically justified in ACS treatment. Adhesion and activation of platelets after erosion and rupture of the atherosclerosis plaque plays a key role in the initialization and development of atherothrombosis.³² Activation of coagulation usually contributes less to the development of acute arterial thrombosis in comparison to antiplatelet action. Rivaroxaban, a direct inhibitor of Xa-factor, does not directly affect the antiplatelet aggregation, caused by collagen, ADP, thromboxane A2 and thrombin.^{33,34} At the same time, it was shown that rivaroxaban, which is capable of total inhibition of thrombin formation, may suppress antiplatelet aggregation inhibiting thrombin generation, through the related tissue factor.³⁵ The main part of thrombin is generated after the primary clot formation, which explains that thrombin plays an important role in clot stabilization rather than its initialization at an early stage of atherothrombosis.³⁶ In previous studies, increased procoagulant activity at an acute stage of ACS was revealed with a significant increase of thrombin formation, which can remain for several months after the event increasing the risk of thrombotic complications.^{37–40} The analysis of randomized clinical trials of the third phase and postmarket studies have shown that high concentrations of DOACs in the blood plasma correlate with the increased rate of bleeding events.^{41,42} Similar studies have shown the connection between low plasma concentrations of DOACs measured in the first month of therapy and

thromboembolic events.^{43,44} In addition, the data from the centers have shown that a 15-fold variation of plasma concentration of rivaroxaban can be observed with an average level of 40 ng/mL. And almost 40% of the patients were beyond the therapeutic range.⁴⁵ In the development of such a variability of pharmacokinetic parameters pharmacogenetics can play a significant role. Rivaroxaban is a substrate of transmembrane transporter P-glycoprotein and is metabolized by P450 enzymes (CYP3A and CYP2J2).⁴⁶ Consequently, the carriership of gene polymorphisms which encode the P-glycoprotein formation (*ABCB1*) and isoenzyme CYP3A4 and CYP3A5 may exert influence on pharmacokinetic, pharmacodynamic parameters and clinical outcomes on rivaroxaban therapy.⁴⁷

That is why the search for safety and effectiveness biomarkers of a combined antithrombotic therapy is of particular interest, and the aim of this study is to assess the influence of gene *CYP2C19*, *CYP3A4*, *CYP3A5* and *ABCB1* polymorphisms on primary and secondary clinical outcomes among patients with ACS and non-valvular AF who received clopidogrel and rivaroxaban.

Materials and Methods

Design and Study Population

A multicenter prospective registry trial was performed to study the efficacy and safety of a combined antithrombotic therapy: 103 patients with non-valvular atrial fibrillation some undergoing a percutaneous coronary intervention were consecutively involved.

The study protocol was approved by the Ethics Committee of Russian Medical Academy of Continuous Professional Education, Moscow, Russian Federation (resolution no. KM-634), it was conducted in accordance with the Declaration of Helsinki, and all patients gave written informed consent for participation.

Inclusion criteria: presence of an informed written consent, age above 18 years, ACS event less than 7 days, non-valvular atrial fibrillation, registered, and which needs anticoagulant treatment (≥ 2 CHA₂DS₂-VASC score for men and ≥ 3 CHA₂DS₂-VASC score for women); first registered atrial fibrillation which needs anticoagulant prescription. Exclusion criteria: pregnancy, lactation, active internal bleeding, liver cirrhosis with C Child-Pugh liver insufficiency; chronic renal disease, HIV-infection, alcohol/drug addiction; mitral valve stenosis moderate and severe, mechanical heart valves, severe psychiatric

disorders, allergic reactions/drug intolerance. The follow-up period was 12 months with telephone interviews every 3 months. The trial assessed the primary outcomes (major bleeding, in-hospital death, cardiovascular death, stroke \transient ischaemic attack, death/renal insufficiency) and secondary outcomes (platelet reactivity units, rivaroxaban concentration)

Assessment of Platelet Activity

The assessment of residual platelet reactivity was performed utilizing VerifyNow assay («Accumetrics», US). For assessment of platelet activity venous blood was used, drained 7 days after the beginning of clopidogrel therapy in vacuum vials 2 mL with 3.2% sodium citrate. Platelets are activated by ADP and interact with fibrinogen on the microparticles in the solution. The more the aggregation is - the less is the optical density of the solution. The level of aggregation of platelets is measured in P2Y₁₂ Reactivity Units or percentage of inhibition. The study was conducted within 1 hr (it is permissible to conduct the study within 4 hrs) after the whole venous blood sample was drained. The therapeutic range of P2Y₁₂-inhibitors when measured with VerifyNow P2Y₁₂ assay are as follows: PRU >208 – the risk of thrombotic events, PRU <95 – bleeding risk, 95 $<$ PRU <208 – adequate response to clopidogrel.

Measurement of Rivaroxaban Plasma Concentration

The blood plasma sampling utilizing VACUETTE[®] (Greiner Bio-One, Austria) with heparin sodium was performed to assess minimal equilibrium concentration of rivaroxaban in the blood plasma before a regular dose of rivaroxaban on the 5th–6th day of the initiation of the therapy. Rivaroxaban concentration determination was carried out by high-performance liquid chromatography with mass-spectrometry detection (HPLC-MS). Agilent 1200 HPLC – system consists of 4-channel pump, mobile phase degasser and chromatographic columns thermostat. The chromatographic separation was done using Agilent Extend-C18 (2.1*100 mm, 3.5 μ m) column at temperature 40°C. The mobile phase: Solution “A” (50 mL 0.1 M solution ammonium acetate and 5 mL formic acid were diluted with unionized water up to 1 liter) and solution “B” (50 mL 0.1 M solution ammonium acetate and 5 mL formic acid were diluted with acetonitrile up to 1 liter). Chromatographic separation was performed in the

isocratic elution mode with the components ratio “A”：“B” = 70:30. The velocity of the chromatographic mobile phase stream was 0.3 mL/min. The volume of the sample was 10 μ L. The analysis was performed within 7 mins. Agilent Triple Quad LC/MS 6410 mass-spectrometer was used (triple quadrupole type) with electrospray ionization in the positive ionization mode. The detection of rivaroxaban mass-spectra was performed in the multiple reaction monitoring mode. The nebulizer gas pressure was 35 psi. The volume velocity of the dry gas flow was 11 L/min, desolvation temperature was 350°C. The value of fragmentation voltage was 135 V, the collision cell voltage was 25 V. Sample preparation was performed using plasma protein precipitation method. The blood plasma samples thawed at room temperature. Then 100 μ L of plasma was placed in Eppendorf vials, 250 μ L of methanol was added, as well as 0.1% hydrogen chloride 9:1 mixture, mixed with Vortex, left alone for 10 mins and mixed once again. Then the acquired samples were centrifuged at 9000 RCF for 10 mins. The supernatant was transferred into chromatographic vials and placed on autosampler. The technique was validated in the 5–1000 ng/mL concentration range. Precision and accuracy values within-run and between runs did not exceed 15% CV (lower limit of quantitation values were lower 20% CV). The samples were stable after freeze-thaw cycle and a month of storage.

Genotyping

Venous blood collected into vacuum vials VACUETTE[®] (Greiner Bio-One, Austria) with ethylene diamine tetra acetate (EDTA) was used for genotyping. Utilizing real-time polymerase chain reaction with commercially available assays («Sintol», Russia; Thermo Fisher Scientific, USA) utilizing Real-Time CFX96 Touch amplifier (Bio-Rad Laboratories, Inc., USA), the carriership of *CYP2C19*, *CYP3A4*, *CYP3A5* and *ABCB1* polymorphisms was detected.

Statistics

Data analysis was carried out in the statistical package IBM SPSS Statistics 23.0. All quantitative variables were tested for normal distribution by the Shapiro–Wilk criterion, resulting in abnormal data distribution ($Z < 1.0$; $p < 0.0001$). For the subsequent analysis of quantitative variables between subgroups, nonparametric criteria (Mann–Whitney, Kruskal–Wallis) were applied. Comparison of frequencies of categorical variables was carried out by means of Pearson’s Chi-square. In order to establish the predictive

role of genetic polymorphisms and clinical parameters for the outcomes, binomial logistic regression was carried out. For the regression analysis, the following variables were selected as the dependent variables: serious bleeding, in-hospital death, death by renal failure, stroke, cardiovascular death. The regression model always included genetic polymorphisms *CYP3A5*3*, *CYP2C19*2*, **17*, *ABCB1* and *DAPT/TATT* variables as covariates. Parameters which showed significant differences when comparing subgroups by outcomes were added as covariates in relevant regression model. Covariates were included in the model using the Wald reverse step-by-step selection method. There were no differences from Hardy-Weinberg equilibrium (Table 1).

Results

The baseline characteristics of the population are featured in Table 2. In the studied patient group 54.5% male, mean age 73 \pm 9.8. PCI, CABG (Coronary artery bypass graft), Myocardial infarction, Heart Failure and bleeding were observed among 32.0%, 6.8%, 46.6%, 35.9% and 19.4%, respectively. Among the major adverse cardiovascular events during the 12-months observation period major bleeding – 11 (10.6%), cardiovascular death – 4 (3.9%), acute cerebrovascular event – 2 (2.0%). The number of patients with resistance to clopidogrel treatment (PRU >208) and who have the risk of bleeding events (PRU <95): 12 (11.7%) and 25 (24.3%), respectively.

Table 3 features the results of the assessment of the influence of polymorphism carriership *CYP2C19*, *CYP3A4*, *CYP3A5* and *ABCB1* among patients with ACS and non-valvular AF on secondary (clopidogrel antiplatelet activity), rivaroxaban equilibrium concentration clinical outcomes. When comparability assessment was performed between the groups of patients, statistically significant differences in terms of factors which could affect the primary and secondary outcomes were observed for *ABCB1 3435 C > T* (T-carriers had higher HAS-BLED, CHA2DS2VASc scores and mean age) and *CYP2C19*2* (*2-carriers had lower Killip scores).

Among the allelic variant carriers *CYP2C19*17* (*CYP2C19*17 CT* + *CYP2C19*17 TT*) patients had a twice more often increased risk of bleeding: low platelet reactivity (PRU <95) was observed among 17.2% of carriers and 33.3% among non-carriers (OR 0.42 95% CI 0.17–1.05, $p = 0.059$). In addition, this allelic variant was significantly associated with the risk of in-hospital death (0% vs 6.7%; OR 0.42 95% CI 0.33–0.53 $p = 0.046$) and cardiovascular death (0% vs 8.9%; OR 0.41

Table 1 Baseline Demographics and Clinical Characteristics (n=103)

Characteristics	Value
Age, years mean \pm SD	73 \pm 9.8
Men, n (%)	56 (54.5%)
Obesity	51 (49.5)
Smoking	9 (8.7%)
CRUSADE Score	11.4 \pm 5.1
Killip Score	1.43 \pm 0.74
CHA2DS2VASc Score	5.1 \pm 1.5
HAS-BLED Score	2.4 \pm 0.9
GFR	53.8 \pm 17.7
Previous PCI	33 (32.0%)
Previous CABG	7 (6.8%)
Previous Myocardial Infarction	48 (46.6%)
Heart Failure	37 (35.9%)
Any Bleeding Events	20 (19.4%)
PCI:	76 (73.8%)
1. Angioplasty	3 (3.9%)
2. Drug-Eluting Stent (DES)	41 (53.9%)
3. Bare-Metal Stent (BMS)	32 (42.1%)
AF:	103 (100%)
1. First Detected	13 (12.6%)
2. Paroxysmal	53 (51.5%)
3. Permanent	37 (35.9%)
MI:	103 (100%)
1. STEMI	28 (27.2%)
2. NSTEMI	52 (50.5%)
3. Unstable angina	23 (22.3%)
Major Bleeding	11 (10.6%)
Cardiovascular Death	4 (3.9%)
Cerebrovascular Event	2 (2.0%)
PRU:	
1. PRU >208	12 (11.7%)
2. PRU <95	25 (24.3%)
Combined antithrombotic therapy	
1. Triple	76 (73.7%)
2. Double	27 (26.3%)

95% CI 0.33–0.52 $p = 0.021$). The *CYP2C19*17* polymorphism carriers had a tendency towards an increase in the frequency of strokes/transient ischaemic attacks (0.0% vs 4.8% OR 0.41 95% CI 0.32–0.52 $p = 0.093$). Patients with T allelic variant polymorphism *ABCB1 rs4148738* and G allelic variant polymorphism *CYP2C19 * 2* had a tendency towards death by renal insufficiency (15.8% vs 35.7% OR 2.96 95% CI 0.80–11.0, $p = 0.093$ and 37.5% vs 19.4% OR 0.40 95% CI 0.15–1.10, $p = 0.07$, respectively). *CYP3A5*3 AG* patients had a tendency towards the increase in the number of major bleeding

Table 2 Hardy-Weinberg Equilibrium Test

Genetic Polymorphism	Genotype	n	%	Chi-Square
<i>CYP3A4*22 C>T intron 6 rs35599367</i>	CC	102	96.2	0
	CT	1	3.8	
<i>CYP3A5*3 rs776746</i>	GG	84	81.6	1.06
	AG	19	18.4	
<i>ABCB1 rs4148738</i>	CC	19	18.4	0.58
	CT	46	44.7	
	TT	38	36.9	
<i>ABCB1 rs1045642</i>	CC	30	29.1	1.63
	CT	45	43.7	
	TT	28	27.2	
<i>CYP2C19*2 rs4244285</i>	AA	2	1.9	0.22
	GA	29	28.2	
	GG	72	69.9	
<i>CYP2C19*17 rs12248560</i>	CC	58	56.3	0.33
	CT	40	38.8	
	TT	5	4.9	

with double/triple antithrombotic therapy: 22.2% vs 8.5% OR 0.33 95% CI 0.08–1.27, $p = 0.093$.

The mean platelet reactivity unit values of patients with T allelic variant polymorphism *ABCB1 rs4148738* were significantly higher compared to non-carriers: 113.3 \pm 57.1 vs 141.0 \pm 62.3 ($p = 0.053$). Carriers of the allelic variant *CYP2C19 * 17 rs12248560* had higher platelet inhibition values (37.8% \pm 28.9 compared to 25.0% \pm 25.9; $p = 0.013$) and, therefore, lower PRU values (121.4 \pm 63.5 vs 147.1 \pm 59.0; $p = 0.044$) compared to non-carriers. *CYP3A4*22 rs35599367* polymorphism had only one heterozygous variant, so, statistic analysis was not performed.

Additional analysis also revealed that higher platelet inhibition level is connected to higher in-hospital death (63.3% vs 29.6%; $p = 0.058$), and higher rivaroxaban concentration is associated with higher HAS-BLED scores: 92.3 \pm 64.9 if HAS-BLED 3–9 vs 67.2 \pm 47.4 if HAS-BLED <3 ($p = 0.03$). Besides, the correlation analysis revealed a weak negative correlation between haemoglobin levels and rivaroxaban plasma concentration ($r=0.3$; $p=0.002$) and platelet activity (0.2; $p=0.02$).

Linkage Disequilibrium Analysis and Haplotype Analysis

We used “SNPStats” for the analysis of linkage disequilibrium and haplotype analysis. The polymorphisms are considered to be in ‘strong LD’ if the one-sided upper

Table 3 Association of Gene Polymorphisms *CYP2C19*, *CYP3A4*, *CYP3A5* and *ABCB1* with Secondary Outcomes

<i>ABCB1</i> 3435 C>T	CC			CT+TT			p-value (Mann-Whitney Test)
	N	M	SD	N	M	SD	
Platelet inhibition rate (%)	30	29.8	28.2	73	30.9	27.9	0.839
Platelet reactivity units (PRU)	30	136.8	61.3	73	135.5	32.8	0.910
Rivaroxaban concentration	30	74.7	54.0	73	80.6	59.0	0.596
<i>ABCB1</i> rs4148738	CC			CT+TT			p
	N	M	SD	N	M	SD	
Platelet inhibition rate (%)	19	34.5	28.5	84	29.7	27.8	0.416
Platelet reactivity units (PRU)	19	113.3	57.1	84	141.0	62.3	0.053
Rivaroxaban concentration	19	78.8	59.2	84	78.9	57.2	0.865
<i>CYP2C19</i> *2	GG			GA+AA			p
	N	M	SD	N	M	SD	
Platelet inhibition rate (%)	72	31.4	26.3	31	28.6	31.6	0.275
Platelet reactivity units (PRU)	72	131.9	57.8	31	145.2	71.0	0.276
Rivaroxaban concentration	72	75.7	57.7	31	86.2	56.9	0.219
<i>CYP2C19</i> *17	CC			CT+TT			p
	N	M	SD	N	M	SD	
Platelet inhibition rate (%)	58	25.0	25.9	45	37.8	28.9	0.013
Platelet reactivity units (PRU)	58	147.1	59.0	45	121.4	63.5	0.044
Rivaroxaban concentration	58	83.5	66.7	45	72.9	42.4	0.821
<i>CYP3A5</i> *3	GG			AG+AA			p
	N	M	SD	N	M	SD	
Platelet inhibition rate (%)	84	30.6	28.0	19	30.4	28.2	0.993
Platelet reactivity units (PRU)	84	135.0	62.9	19	139.8	60.0	0.75
Rivaroxaban concentration	84	81.6	58.9	19	69.7	55.9	0.430

95% confidence bound on D' is ≥ 0.98 and the lower bound is ≥ 0.70 .

The polymorphisms in the *CYP2C19* gene (*2 (*rs4244285*) and *17 (*rs12248560*)) were in high linkage disequilibrium ($D' = 0.99$). There was no AA genotype among the patients who had the bleeding. Major bleeding showed a significant association ($p < 0.05$) with

a protective effect of *CYP2C19**2 *rs4244285* in the presence of *CYP2C19**17 *rs12248560* against major bleeding (Table 4)

Discussion

Although the optimal values of therapeutic range of rivaroxaban concentration in blood plasma have not been

Table 4 Haplotype Analysis of the *CYP2C19* Polymorphisms for Major Bleeding

Haplotype Association with Response (n=103, Crude Analysis)					
	<i>CYP2C19</i> *2	<i>CYP2C19</i> *17	Freq	OR (95% CI)	P-value
1	G	C	0.6101	1.00	–
2	G	T	0.2297	0.13 (0.03–0.60)	0.01
3	A	C	0.1472	0.00 (-Inf - Inf)	1
Global haplotype association p-value: 0.0046					

determined, in previous studies significant associations between equilibrium rivaroxaban concentration and incidence of bleeding among AF patients were revealed (48 ± 30 vs 34 ± 26 ; $p = 0.02$).⁴⁸ In addition, apart from acknowledged factors (age, renal insufficiency^{49,50}), genetic factors such as gene polymorphisms *ABCB1*, *CYP3A4* may exert influence on rivaroxaban pharmacokinetics and, therefore, on the risk of bleeding complications. This is extremely important in case of antithrombotic therapy among patients with AF and ACS, even more so in light of the correlation of the aforementioned genetic markers with pharmacologic response to clopidogrel,² while rivaroxaban indirectly affects platelet aggregation through thrombin generation inhibition.³⁵ Previous studies assessed primarily non-genetic risk factors of poor outcomes among patients with AF and ACS. Less emphasis was put on the role of genetic markers. Zang L. et al associated major bleeding with rivaroxaban concentration (2.5 mg), sex and prior myocardial revascularization.⁵¹ Several studies assessed the role of genetic factors among patients with AF without ACS. Anne-Laure Sennesael et al report that three patients out of four having major bleeding and blood rivaroxaban concentration above expected (more than 136 nm/mL) were heterozygous *ABCB1 1236C > T*, *2677G > T*, *3435 C > T* and *rs4148738*.²⁰ Likewise, *ABCB1 2677G > T* and *3435 C > T* carriership were associated with higher blood plasma rivaroxaban concentration and bleeding complications.⁵² Apart from that, there are studies that report no link between *1236C > T*, *2677G > T*, *3435 C > T* and rivaroxaban pharmacokinetics.⁵³

The present study has proven that gene polymorphisms, usually associated with clopidogrel-related bleeding, play an important role in combined antithrombotic therapy: *CYP2C19*17* carriers twice as often had low residual platelet reactivity (PRU less than 95) 17.2% vs 33.3% ($p = 0.059$).

*CYP2C19*17* carriers turned out to be prone to increased platelet inhibition ($37.8\% \pm 28.9$ compared to $25.0\% \pm 25.9$; $p = 0.013$) and, therefore, lower PRU values (121.4 ± 63.5 vs 147.1 ± 59.0 ; $p = 0.044$) compared with non-carriers.

Apart from that, this allelic variant turned out to be the only genetic marker to be significantly associated with in-hospital (0% vs 6.7%; $p = 0.046$) and cardiovascular death (0% vs 8.9%; $p = 0.021$).

No genetic marker was connected to clopidogrel resistance (PRU more than 208). The only allelic variant which led towards higher mean PRU values turned out to be *ABCB1 rs4148738*: 113.3 ± 57.1 vs 141.0 ± 62.3 ($p = 0.053$).

The results can be explained by the fact that in situations of indirect influence of rivaroxaban on platelet reactivity, it is safe to put more emphasis on the risk factors of bleeding complications rather than on well described in literature genetic risk factors for thrombotic complications on clopidogrel therapy (low function alleles *CYP2C19*2 rs4244285*, *CYP2C19*3 rs4986893*), as these bleeding risk factors do not demand a switch of drug according to the present CPIC guidelines and to a lesser degree disrupt clopidogrel response.²

Conclusion

*CYP2C19*17 rs12248560* is significantly associated with in-hospital death of patients with an acute coronary syndrome and non-valvular atrial fibrillation who receive clopidogrel and rivaroxaban.

None of the markers under study (*CYP3A5*3 rs776746*, *CYP2C19*2 rs4244285*, **17 rs12248560*, *ABCB1 3435 C > T*, *ABCB1 rs4148738*) has proven to affect rivaroxaban equilibrium concentration in blood plasma among patients with atrial fibrillation and acute coronary syndrome.

In situations of double or triple antithrombotic rivaroxaban and clopidogrel therapy among patients with AF and ACS, the genetic factors associated with bleeding complications risk (*CYP2C19*17 rs12248560*) may prove to be clinically relevant.

Additional results regarding a connection between rivaroxaban concentration and high HAS-BLED scores demonstrate the feasibility of using this marker for personalization of rivaroxaban therapy.

Acknowledgment

This study has been supported by the Russian Science Foundation under Project No. 16-15-00227.

Disclosure

Dmitriy Ivashchenko reports grants from Russian Science Foundation and RFBR, outside the submitted work. The other authors report no conflicts of interest in this work.

References

1. Levine GN, Bates ER, Bittl JA, et al. ACC/AHA guideline focused update on duration of dual antiplatelet therapy in patients with coronary artery disease: a report of the American College of Cardiology/American Heart Association Task Force on clinical practice guidelines. *J Am Coll Cardiol.* 2016;2016(68):1082–1115. doi:10.1016/j.jacc.2016.03.513

2. Scott SA, Sangkuhl K, Stein CM, et al. Clinical pharmacogenetics implementation consortium. Clinical pharmacogenetics implementation consortium guidelines for CYP2C19 genotype and clopidogrel therapy: 2013 update. *Clin Pharmacol Ther.* 2013;94:317–323. doi:10.1038/clpt.2013.105
3. Sibbing D, Koch W, Gebhard D, et al. Cytochrome 2c19*17 allelic variant, platelet aggregation, bleeding events, and stent thrombosis in clopidogrel-treated patients with coronary stent placement. *Circulation.* 2010;121:512–518. doi:10.1161/CIRCULATIONAHA.109.885194
4. Ellis KJ, Stouffer GA, McLeod HL, Lee CR. Clopidogrel pharmacogenomics and risk of inadequate platelet inhibition: US FDA recommendations. *Pharmacogenomics.* 2009;10:1799–1817. doi:10.2217/pgs.09.143
5. O'Donoghue M, Wiviott SD. Clopidogrel response variability and future therapies: clopidogrel: does one size fit all? *Circulation.* 2006;114:e600–e606. doi:10.1161/CIRCULATIONAHA.106.643171
6. Tam CC, Kwok J, Wong A, et al. Genotyping-guided approach versus the conventional approach in selection of oral P2Y12 receptor blockers in Chinese patients suffering from acute coronary syndrome. *J Int Med Res.* 2017;45(1):134–146. doi:10.1177/030060516677190
7. Koltowski L, Tomaniak M, Aradi D, et al. Optimal antiplatelet pharmacotherapy guided by bedSIDE genetic or functional TESTING in elective PCI patients: a pilot study: ONSIDE TEST pilot. *Cardiol J.* 2017;24(3):284–292. doi:10.5603/CJ.a2017.0026
8. So DYF, Wells GA, McPherson R, et al. A prospective randomized evaluation of a pharmacogenomic approach to antiplatelet therapy among patients with ST-elevation myocardial infarction: the RAPID STEMI study. *Pharmacogenomics J.* 2016;16(1):71–78.
9. Roberts JD, Wells GA, Le May MR, et al. Point-of-care genetic testing for personalisation of antiplatelet treatment (RAPID GENE): a prospective, randomised, proof-of-concept trial. *Lancet.* 2012;379(9827):1705–1711. doi:10.1016/S0140-6736(12)60161-5
10. Shen DL, Wang B, Bai J, et al. Clinical value of CYP2C19 genetic testing for guiding the antiplatelet therapy in a Chinese population. *J Cardiovasc Pharmacol.* 2016;67:232–236. doi:10.1097/FJC.0000000000000337
11. Xie X, Ma YT, Yang YN, et al. Personalized antiplatelet therapy according to CYP2C19 genotype after percutaneous coronary intervention: a randomized control trial. *Int J Cardiol.* 2013;168:3736–3740. doi:10.1016/j.ijcard.2013.06.014
12. Notarangelo FM, Maglietta G, Bevilacqua P, et al. Pharmacogenomic approach to selecting antiplatelet therapy in acute coronary syndromes: PHARMCLO trial. *J Am Coll Cardiol.* 2018;71(17):1869–1877. doi:10.1016/j.jacc.2018.02.029
13. Sánchez-Ramos J, Dávila-Fajardo CL, Toledo Frías P, et al. Results of genotype-guided antiplatelet therapy in patients who undergone percutaneous coronary intervention with stent. *Int J Cardiol.* 2016;225:289–295. doi:10.1016/j.ijcard.2016.09.088
14. Deiman BALM, Tonino PAL, Kouhestani K, et al. Reduced number of cardiovascular events and increased cost-effectiveness by genotype-guided antiplatelet therapy in patients undergoing percutaneous coronary interventions in the Netherlands. *Neth Heart J.* 2016;24(10):589–599. doi:10.1007/s12471-016-0873-z
15. Cavallari LH, Franchi F, Rollini F, et al. Clinical implementation of rapid CYP2C19 genotyping to guide antiplatelet therapy after percutaneous coronary intervention. *J Transl Med.* 2018;16(1):92. doi:10.1186/s12967-018-1469-8
16. Klein MD, Lee CR, Stouffer GA. Clinical outcomes of CYP2C19 genotype-guided antiplatelet therapy: existing evidence and future directions. *Pharmacogenomics.* 2018;19:1039–1046. doi:10.2217/pgs-2018-0072
17. Mega JL, Simon T, Collet JP, et al. Reduced-function CYP2C19 genotype and risk of adverse clinical outcomes among patients treated with clopidogrel predominantly for PCI: a meta-analysis. *JAMA.* 2010;304:1821–1830. doi:10.1001/jama.2010.1543
18. Kheiri B, Osman M, Abdalla A, et al. CYP2C19 pharmacogenetics versus standard of care dosing for selecting antiplatelet therapy in patients with coronary artery disease: a meta-analysis of randomized clinical trials. *Catheter Cardiovasc Interv.* 2019;93(7):1246–1252. doi:10.1002/ccd.27949
19. Wallentin L, James S, Storey RF, et al.; PLATO investigators. Effect of CYP2C19 and ABCB1 single nucleotide polymorphisms on outcomes of treatment with ticagrelor versus clopidogrel for acute coronary syndromes: a genetic substudy of the PLATO trial. *Lancet.* 2010;376:1320–1328. doi:10.1016/S0140-6736(10)61274-3
20. Mega JL, Close SL, Wiviott SD, et al. Cytochrome P450 genetic polymorphisms and the response to prasugrel: relationship to pharmacokinetic, pharmacodynamic, and clinical outcomes. *Circulation.* 2009;119:2553–2560. doi:10.1161/CIRCULATIONAHA.109.851949
21. Wiviott SD, Braunwald E, McCabe CH, et al.; TRITON-TIMI 38 Investigators. Prasugrel versus clopidogrel in patients with acute coronary syndromes. *N Engl J Med.* 2007;357:2001–2015. doi:10.1056/NEJMoa0706482
22. Wallentin L, Becker RC, Budaj A; PLATO Investigators, et al. Ticagrelor versus clopidogrel in patients with acute coronary syndromes. *N Engl J Med.* 2009;361:1045–1057. doi:10.1056/NEJMoa0904327
23. Dayoub EJ, Seigerman M, Tuteja S, et al. Trends in platelet adenosine diphosphate P2Y12 receptor inhibitor use and adherence among antiplatelet-naïve patients after percutaneous coronary intervention, 2008–2016. *JAMA Intern Med.* 2018;178:943–950. doi:10.1001/jamainternmed.2018.0783
24. January CT, Wann LS, Calkins H, et al. 2019 AHA/ACC/HRS focused update of the 2014 AHA/ACC/HRS guideline for the management of patients with atrial fibrillation: a report of the American College of Cardiology/American Heart Association task force on clinical practice guidelines and the heart rhythm society. *J Am Coll Cardiol.* 2019;74(1):104–132. doi:10.1016/j.jacc.2019.01.011
25. Mega JL, Braunwald E, Mohanavelu S, et al. Rivaroxaban versus placebo in patients with acute coronary syndromes (ATLAS ACS-TIMI 46): a randomised, double-blind, Phase II trial. *Lancet.* 2009;374:29–38. doi:10.1016/S0140-6736(09)60738-8
26. Oldgren J, Budaj A, Granger CB, et al. Dabigatran vs. placebo in patients with acute coronary syndromes on dual antiplatelet therapy: a randomized, double-blind, phase II trial. *Eur Heart J.* 2011;32:2781–2789. doi:10.1093/eurheartj/ehr113
27. Alexander JH, Becker RC, Bhatt DL, et al. Apixaban, an oral, direct, selective factor Xa inhibitor, in combination with antiplatelet therapy after acute coronary syndrome: results of the Apixaban for Prevention of Acute Ischemic and Safety Events (APPRAISE) trial. *Circulation.* 2009;119:2877–2885.
28. Gibson CM, Mehran R, Bode C, et al. Prevention of bleeding in patients with atrial fibrillation undergoing PCI. *N Engl J Med.* 2016;375:2423–2434. doi:10.1056/NEJMoa1611594
29. Cannon CP, Bhatt DL, Oldgren J, et al. Dual antithrombotic therapy with dabigatran after PCI in atrial fibrillation. *N Engl J Med.* 2017;377:1513–1524. doi:10.1056/NEJMoa1708454
30. Cavallari I, Patti G. Meta-analysis comparing the safety and efficacy of dual versus triple antithrombotic therapy in patients with atrial fibrillation undergoing percutaneous coronary intervention. *Am J Cardiol.* 2018;121:718–724. doi:10.1016/j.amjcard.2017.12.014
31. Golwala HB, Cannon CP, Steg PG, et al. Safety and efficacy of dual vs. triple antithrombotic therapy in patients with atrial fibrillation following percutaneous coronary intervention: a systematic review and meta-analysis of randomized clinical trials. *Eur Heart J.* 2018;39:1726–1735a. doi:10.1093/eurheartj/ehy162
32. Borissoff JI, Spronk HM, Ten Cate H. The hemostatic system as a modulator of atherosclerosis. *N Engl J Med.* 2011;364:1746–1760. doi:10.1056/NEJMra1011670
33. Perzborn E, Strassburger J, Wilmen A, et al. Biochemical and pharmacologic properties of BAY 59-7939, an oral, direct factor Xa inhibitor. *Pathophysiol Haemost Thromb.* 2004;33:Abstract PO079.

34. Fareed J, Hoppensteadt D, Maddenini J, et al. Antithrombotic mechanism of action of BAY 59-7939 – a novel, oral, direct factor Xa inhibitor. *J Thromb Haemost.* 2005;3:Abstract P0518.
35. Wong PC, Jiang X. Apixaban, a direct factor Xa inhibitor, inhibits tissue-factor induced human platelet aggregation in vitro: comparison with direct inhibitors of factor VIIa, XIa and thrombin. *Thromb Haemost.* 2010;104(2):302–310. doi:10.1160/TH10-02-0097
36. Mann KG, Brummel K, Butenas S. What is all that thrombin for? *J Thromb Haemost.* 2003;1:1504–1514. doi:10.1046/j.1538-7836.2003.00298.x
37. Ardissino D, Merlini PA, Bauer KA, et al. Coagulation activation and long-term outcome in acute coronary syndromes. *Blood.* 2003;102:2731–2735. doi:10.1182/blood-2002-03-0954
38. Undas A, Szuldrzyński K, Brummel-Ziedins KE, Tracz W, Zmudka K, Mann KG. Systemic blood coagulation activation in acute coronary syndromes. *Blood.* 2009;113:2070–2078. doi:10.1182/blood-2008-07-167411
39. Ten Cate H, Hemker HC. Thrombin generation and atherothrombosis: what does the evidence indicate? *J Am Heart Assoc.* 2016;5:e003553. doi:10.1161/JAHA.116.003553
40. Merlini PA, Ardissino D, Bauer KA, et al. Persistent thrombin generation during heparin therapy in patients with acute coronary syndromes. *Arterioscler Thromb Vasc Biol.* 1997;17:1325–1330. doi:10.1161/01.ATV.17.7.1325
41. Reilly PA, Lehr T, Haertter S, et al. The effect of dabigatran plasma concentrations and patient characteristics on the frequency of ischemic stroke and major bleeding in atrial fibrillation patients: the RE-LY trial (randomized evaluation of long-term anticoagulation therapy). *J Am Coll Cardiol.* 2014;63(4):321–328. doi:10.1016/j.jacc.2013.07.104
42. Southworth MR, Reichman ME, Unger EF. Dabigatran and postmarketing reports of bleeding. *N Engl J Med.* 2013;368(14):1272–1274. doi:10.1056/NEJMp1302834
43. Testa S, Paoletti O, Legnani C, et al. Low drug levels and thrombotic complications in high-risk atrial fibrillation patients treated with direct oral anticoagulants. *J Thromb Haemost.* 2018;16:842–848. doi:10.1111/jth.2018.16.issue-5
44. Larsen TB, Rasmussen LH, Skjøth F, et al. Efficacy and safety of dabigatran etexilate and warfarin in “real-world” patients with atrial fibrillation: a prospective nationwide cohort study. *J Am Coll Cardiol.* 2013;61(22):2264–2273. doi:10.1016/j.jacc.2013.03.020
45. Seiffge DJ, Traenka C, Polymeris A, et al. Feasibility of rapid measurement of Rivaroxaban plasma levels in patients with acute stroke. *J Thromb Thrombolysis.* 2017;43(1):112–116. doi:10.1007/s11239-016-1431-7
46. Cavallari LH, Shin J, Perera MA. Role of pharmacogenomics in the management of traditional and novel oral anticoagulants. *Pharmacotherapy.* 2011;31:1192–1207. doi:10.1592/phco.31.12.1192
47. Sennesael AL, Larock AS, Douxfils J, et al. Rivaroxaban plasma levels in patients admitted for bleeding events: insights from a prospective study. *Thromb J.* 2018;12(16):28. doi:10.1186/s12959-018-0183-3
48. Miklič M, Mavri A, Vene N, et al. Intra- and inter-individual rivaroxaban concentrations and potential bleeding risk in patients with atrial fibrillation. *Eur J Clin Pharmacol.* 2019;75(8):1069–1075. doi:10.1007/s00228-019-02693-2
49. Akhtar T, Mattumpuram J, Fugar S, et al. Factors associated with bleeding events in patients on rivaroxaban for nonvalvular atrial fibrillation: a real world experience. *JACC.* 2019;73(9). doi:10.1016/S0735-1097(19)31079-4
50. Beyer-Westendorf J, Forster K, Pannach S, et al. Rates, management, and outcome of rivaroxaban bleeding in daily care: results from the Dresden NOAC registry. *Blood.* 2014;124:955–962. doi:10.1182/blood-2014-03-563577
51. Zhang L, Yan X, Nandy P, et al. Influence of model-predicted rivaroxaban exposure and patient characteristics on efficacy and safety outcomes in patients with acute coronary syndrome. *Ther Adv Cardiovasc Dis.* 2019;13:1–15. doi:10.1177/1753944719863641
52. Ing Lorenzini K, Daali Y, Fontana P, Desmeules J, Samer C. Rivaroxaban-induced hemorrhage associated with ABCB1 genetic defect. *Front Pharmacol.* 2016;7:494. doi:10.3389/fphar.2016.00494
53. Gouin-Thibault I, Delavenne X, Blanchard A, et al. Interindividual variability in dabigatran and rivaroxaban exposure: contribution of ABCB1 genetic polymorphisms and interaction with clarithromycin. *J Thromb Haemost.* 2017;15:273–283. doi:10.1111/jth.13577

Pharmacogenomics and Personalized Medicine

Publish your work in this journal

Pharmacogenomics and Personalized Medicine is an international, peer-reviewed, open access journal characterizing the influence of genotype on pharmacology leading to the development of personalized treatment programs and individualized drug selection for improved safety, efficacy and sustainability. This journal is indexed

on the American Chemical Society's Chemical Abstracts Service (CAS). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/pharmacogenomics-and-personalized-medicine-journal>

Dovepress