ADME pharmacogenetics: future outlook for Russia

Karin B Mirzaev¹, Denis S Fedorinov^{*, 1}, Dmitry V Ivashchenko¹ & Dmitry A Sychev¹ ¹Russian Medical Academy of Continuous Professional Education of the Ministry of Healthcare, Moscow, Russian Federation *Author for correspondence: fedorinov.denis@gmail.com

This systematic review reflects the results of pharmacogenetic studies in the Russian Federation aimed at studying the genes involved in the drug biotransformation system. The works of Russian researchers found by us are mostly devoted to microsomal liver oxidation enzymes (metabolism) and membrane transporter systems (absorption and excretion). This review presents population-ethnic and associative clinical studies on the genes of the *CYP450* system, noncytochrome oxidation enzymes (*SULT1A1, CES1*), membrane transporter system genes (*ABCB1, SLCO1B1*) and warfarin biotransformation enzymes (*VKORC1, GGCX*). The information is structured in the form of 11 tables, divided by regions of the Russian Federation.

First draft submitted: 7 February 2019; Accepted for publication: 1 April 2019; Published online: 1 August 2019

Keywords: ADME • cardiology • P450 • personalized medicine • pharmacogenomics • Russia


In recent years, growing attention has been paid to the principles of personalized medicine in the treatment of many diseases in the Russian Federation. Although Hippocrates pointed to the importance of these principles [1]: "It is more important to know what sort of person has a disease than to know what sort of disease a person has" – their application is only now becoming possible. Fundamental knowledge accumulated during the development of evidence-based medicine showed that the genotypic and phenotypic characteristics of the patient determine the predisposition to the disease, response to the drug and effectiveness of treatment in general [2]. The use of pharmacogenetics plays an important role in eliminating the variability of the response to a drug. During preclinical studies, the main metabolic pathway of metabolism is established for each drug. Further on, it is necessary to conduct a large number of additional studies to clarify specific stages of metabolism. Identified application points in the form of transporters, receptors, enzymes, ion channels – among others, may differ by their functional capabilities, which is associated with changes in the genes that encode the listed targets. In turn, the determination of these genetic features of the patient allows prediction of response to the drug.

The Clinical Pharmacogenetics Implementation Consortium (CPIC[®]) accumulates all the knowledge in this field. To date, they have developed complete recommendations for dosing or prioritizing 35 drugs, taking into account the results of pharmacogenetic testing and 335 proven genes associations [3]. For a more convenient presentation of the results of pharmacogenetic testing, CPIC proposed the division of patients into phenotypic groups: normal metabolizer, rapid metabolizer, ultrarapid metabolizer, intermediate metabolizer, poor metabolizer. According to this division, further recommendations for dosing or priority selection of medicines are based.

With the accumulation of knowledge in pharmacogenetics, racial and ethnic particularities in the prevalence of genetic polymorphisms have begun to emerge. At the same time, the carriage frequency of allelic variants may vary significantly among representatives of different races and nationalities. This becomes especially important for such multinational countries as the Russian Federation, which numbers 194 different ethnic groups [4].

The objective of this systematic review was to summarize current research in the field of pharmacogenetics in the Russian Federation.

Future Medicine






Figure 1. Flowchart of the literature selection process.

Materials & methods

Search strategy

The authors conducted an online search for articles in December 2018 using the following databases: PubMed, Google Scholar, eLIBRARY (Russian electronic library of scientific publications). The search strategy combined key terms «Russia», «Russian» with terms associated with polymorphic marker including: «P450», «CYP2C19», «CYP2D6», «CYP2D6», «CYP2B1», «CYP2B6», «CYP2C8», «CYP2C9», «CYP3A4», «CYP3A5», «CYP1A1», «CYP1A2», «CYP4F2», «CYP4F1», «ABCB1», «SLCO1B1», «VKORC1», «GGCX», «SULT1A1», «CES1», «gene», «genes», «pharmacogenetics», «pharmacogenomics», «ethnic group». A thorough review of the references in relevant articles was also completed with citations from all searches.

Systematic review

The search produced 443 articles (Figure 1). After removing duplicates, the titles and abstracts were screened for relevance based on inclusion criteria – the presence of results of pharmacogenetic testing. A total of 369 articles were excluded due to non-relevance of the inclusion criteria. After analyzing 74 full-text articles, 18 were excluded due to the inability to reproduce the results of genotyping and to calculate the prevalence rate of the alleles.

Results & discussion

All processes associated with the pharmacokinetics of a substance in the body are described by the abbreviation ADME, where A is absorption, D is distribution, M is metabolism and E is excretion. The work of Russian researchers collated by us are mostly devoted to microsomal liver oxidation enzymes (metabolism) and membrane transporter systems (absorption and excretion). Within the framework of this systematic review, two groups were identified: clinical trials involving healthy volunteers (designated as healthy) and clinical trials in patients with a specific nosology. If the authors of the study did not indicate the ethnicity of the research participants, we marked this characteristic as 'general population'. In addition, we identified three categories of research: purely population-ethnic works, which studied only the carriage frequency of allelic variants among representatives of different ethnic groups, associative clinical studies that investigated the relationship between the carrier state of allelic variants and the change in a certain physiological parameter of research participants as well as work aimed at developing drug dosing algorithms, taking into account the results of pharmacogenetic testing. Next, Tables 1–10 will present population-ethnic data, and the subsections describe the results of associative studies. Data on the study of minor alleles are presented in Supplementary Table 1.

Region of the	Ethnic group	Number of	Characteristics of the			Allel	ic variants			Re
Russian Federation		patients	population		*2			*3		
				Wt/Mt	Mt/Mt	%	Wt/Mt	Mt/Mt	%	
Dagestan Republic	Laks	46	Healthy	-	-	-	12	3	19.56	[!
	Avars	90		-	-	-	22	3	15.55	
	Dargins	50		-	-	-	15	1	17.00	
European part of Rus	sia:									
– Moskow	Russians	63	Atrial fibrillation	13	2	13.49	14	2	14.29	[(
	Russians	352	Healthy	80	2	11.90	33	1	5.00	[7
	Russians	83	B-cell chronic lymphocytic leukemia	20	0	12.04	13	0	7.83	[8
		76	Non-Hodgkin lymphoma	9	0	5.92	15	1	11.18	
		177	Healthy	29	0	8.19	21	0	5.93	
	General population	91	Warfarin taking	17	2	11.54	18	1	10.99	[9
– Voronezh	Russians	290	Healthy	57	2	10.51	37	1	6.72	[10
– Saint Petersburg	General population	298	Healthy	58	4	11.07	37	1	6.54	[11
		62	Warfarin taking	11	0	8.87	6	0	4.84	
– Ufa	Russians	319	Healthy	17	0	7.59	24	0	9.02	[1]
	Tatars	279		11	0	5.14	14	0	5.43	
	Bashkirs	144		7	0	6.14	8	0	6.15	
– Karachay- Cherkessia	Circassians	77	Healthy	-	-	-	19	0	12.30	[13
	Karachays	125		_	_	_	58	2	24.8	
– Stavropol	Russians	25	Healthy	6	0	12.00	3	0	6.00	[14
	Armenians	25		7	0	14.00	7	0	14.00	
	Karachays	25		2	0	4.00	5	0	10.00	
	Circassians	32		4	0	6.00	6	0	9.00	
Siberia:										
– Yakutsk	Yakuts	229	Healthy	7	1	5.05	10	1	6.74	[15
– Cheriktei	Yakuts	88	Healthy	2	0	1.10	1	0	0.60	[16
– Kyzil	Tuvinians	88		2	0	1.10	9	0	5.10	
– Buriatia	Buryats	88		4	0	2.30	3	0	1.70	
– Altai	Altaians	87		10	0	5.70	14	1	9.20	
– Tomsk	Russians	87		19	1	12.10	8	1	6.80	
	General population	250	Breast cancer	11	0	8.87	36	3	9.00	[1]
Far East:										
– Khabarovsk	Nanai	70	Healthy	0	0	0.00	8	0	5.70	[18
Ural:										
– Chelyabinsk	General population	63	Cardiac patients	13	1	11.90	10	0	7.94	[19

CYP450 enzymes CYP2C9

Cytochrome *CYP2C9* gene has more than 60 allelic variants [20], but in clinical practice the largest role is played by alleles *2, *3, *5, *6, *8 and *11, the carrier of which is associated with reduced function of the enzyme. CYP2C9 is involved in the metabolism of several important drugs with low therapeutic index, namely: S-warfarin, fluvastatin, S-acenocoumarol, ibuprofen, phenytoin, tolbutamide, nateglinide. Accordingly, it is necessary to take into account carriage of slow alleles of *CYP2C9* gene encoding the same name isozyme, in order to avoid adverse drug effects in the appointment of listed drugs [21]. In particular, carriage of *2, *3, *5 and *6 alleles are taken into account when

Region of the Russian	Ethnic group	Number of	Characteristics of the population		Allelic varia	nts	Ref.
Federation		patients		Wt/Mt	Mt/Mt	%	
CYP2C9 rs9332242							
Kursk	General population	561	CHD	100	3	9.40	[25]
		694	Healthy	124	12	10.70	
		425	Essential hypertension	80	4	10.50	[26]
		391	Healthy	71	7	11.10	
CYP2C9 rs4918758							
Kursk	General population	561	СНД	257	42	30.40	[25]
		694	Healthy	302	82	33.60	
CYP2C9 rs61886769							
Kursk	General population	561	СНД	168	16	17.80	[25]
		694	Healthy	220	23	19.20	
VKORC1 - 1639 G > A rs992	23321						
Yakutsk	Yakuts	89	Healthy	28	60	83.14	[15]
Moscow	Russians	63	Atrial fibrillation	24	12	38.10	[6]
Moscow	General population	91	Warfarin taking	26	20	36.26	[9]
GGCX rs11676382							
Moscow	Russians	63	Atrial fibrillation	6	0	4.76	[6]

calculating the dose of warfarin (available on the resource WarfarinDosing.org) [22]. When identifying carriage of *8 and *11 alleles in patients of African descent, it is additionally recommended to reduce the calculated dose by 15–30% [23].

The carriage frequency of these allelic variants differs among representatives of various ethnic groups. *CYP2C9*2* is most common among Middle Eastern – 13.21%, Europeans – 12.60%, south/central Asian – 10.74%, quite rare among Americas – 6.25% and practically does not occur among representatives of African – 2.36%, African–American – 2.30% and east Asian – 0.06%. *CYP2C9*3* is common among representatives of south and central Asian – 10.17%, Middle Eastern – 9.31%, Caucasian – 7.08% but rarely occurs in the African–American – 1.17% and African population – 1.03%; *5, *6, *8 and *11 allelic variants with different frequencies occur mainly in people of African origin 0.77–6.66% [24].

At present, a large number of pharmacogenetic studies have been conducted on this gene in Russia. Cumulative information about the main characteristics of the studied population and the allele frequencies of the studied gene are presented in Tables 1 and 2.

Analysis of Russian works shows that the prevalence frequency of allelic variants *CYP2C9*2* and *CYP2C9*3* has its own peculiarities. Thus, *2 allelic variant prevails among Russians and Armenians as representatives of the Europoid race, and is rarely found among representatives of the Mongoloid race, namely: Yakuts, Buryats, Tuvinians. What is important, *3 allelic variant is widespread among representatives of the Caucasian peoples: Circassians, Laks, Avars and Dargins – which indicates the validity of pharmacogenetic testing in this group of patients.

The analysis of the results also revealed a specific feature of our country: the frequency of allelic variants prevalence both among Russians and in the general population changes depending on the region. This is probably due to population migration and inter-ethnic marriages.

One of the first applications of pharmacogenetic testing began in the Moscow region. Gra *et al.* studied the carriage frequency of *CYP2C9* allelic variants in Russian patients with B-cell chronic lymphocytic leukemia and Non-Hodgkin lymphoma in comparison with healthy patients. The study revealed a higher prevalence of allelic variant *CYP2C9*2* in men with B-cell chronic lymphocytic leukemia (OR = 2.38; 95% CI: 1.1–5.2), as well as a combination of *CYP2C9*2* and *CYP2C9*3* (OR = 2.09; 95% CI: 1.1–3.9) [8]. The results were used to create a biochip. Ivashchenko *et al.* studied the general population of patients taking warfarin [9]. The obtained results turned out to be comparable with data on the carriage frequency of allelic variants of this gene among representatives of the Caucasians.

Barysheva *et al.* studied the prevalence of *CYP2C9* allelic variants in cardiac patients in the Ural region [19]. Based on the results of pharmacogenetic testing, personalized selection of the warfarin dose was carried out.

Polonikov *et al.* studied rs9332242 and rs4918758 polymorphisms of *CYP2C9* and their possible role in the development of susceptibility to coronary heart disease (CHD) and essential hypertension, but the result was negative [25,26].

Pchelina *et al.* investigated individual sensitivity to warfarin in patients with atrial fibrillation and prosthetic heart valves who received this drug. The results were compared with a healthy population. An association was established between the carriage of *CYP2C9*2* and *CYP2C9*3* alleles and excessive anticoagulation [11].

In Siberia, Seredina *et al.* studied the effectiveness of neoadjuvant chemotherapy in patients with breast cancer [17]. *CYP2C9*2* was associated with neoadjuvant chemotherapy resistance (OR = 4.64; 95% CI: 1.01-20.91).

Sychev *et al.* showed the ineffectiveness of existing Acenocoumarol dosing algorithms based on *CYP2C9* geno-typing for Russian patients with Atrial fibrillation [6].

In addition to cytochrome CYP2C9, other enzymes are involved in the metabolism of Warfarin. γ -Glutamyl carboxylase (*GGCX*) oxidizes vitamin K hydroquinone (the active form of vitamin K1) to vitamin K 2,3 epoxide [27]. VKORC1 returns the epoxides formed in the carboxylation reaction to the active form of vitamin K. It becomes clear that changes in the activity of these enzymes will lead to shifts in the vitamin K metabolism cycle and, as a result, changes in the effectiveness of anticoagulant therapy. Therefore, the polymorphisms of these enzymes are taken into account in the dosing regimen of anticoagulants [28].

Vasilyev *et al.* found a high prevalence of *VKORC1* – 1639G> A rs9923321 polymorphism in the Yakut population – 83.14% [15], while the prevalence among patients of the Moscow region was only 36.26% according to Ivashchenko *et al.* [9]. The work mentioned earlier by Sychev *et al.* did not show the effectiveness of existing Acenocoumarol dosing algorithms with regard to the *VKORC1* genotype as applied to the Russian population [6].

CYP2C19

Cytochrome CYP2C19 enzyme is involved in the metabolism of several important classes of drugs: benzodiazepines [29], antidepressants [30], proton pump inhibitors (PPIs) [31], mephenytoin [32] and the antiplatelet prodrug clopidogrel [33]. Currently, 35 allelic variants of the *CYP2C19* gene have been discovered [34]. In pharmacogenetic testing, three alleles are used as polymorphic markers: *2, *3, encoding an enzyme with reduced activity, and *17, a change in the promoter zone, resulting in the formation of a larger amount of enzyme. It should be noted that the simultaneous presence of these two mutations is possible. In this case, the person will have increased expression of the enzyme with reduced activity [35].

The carrier rate of *CYP2C19*2* allelic variant can reach 53.9% among representatives of the Oceanian ethnic groups, common among south/central Asian – 34.3%, east Asian – 29% and is quite rare among representatives of the African–American – 18.3%, Europeans – 14.6% and Middle Eastern – 13.2%. *CYP2C19*3* allelic variant of is not so widespread in the world population: the carriage frequency among representatives of Oceania's ethnic groups reaches 14.5%, among Asians 4–5%, and is quite rare among other ethnic groups – 0.28–0.73%. The reverse trend is observed for *CYP2C19*17* allelic variant: it is rarely found in Oceania - 4.1% and east Asian – 1.31%, but is widespread in Middle Eastern – 21.6%, Caucasians – 21.3% and African–American – 21% [36].

Cumulative information about the main characteristics of the studied population and the frequency of *CYP2C19* alleles are presented in Table 3.

Analysis of the work of Russian researchers revealed a wide spread of *CYP2C19*2* and *CYP2C19*17* allelic variants in the population, while *CYP2C19*3* is less common and is mainly found in representatives of the Mongoloid race. The prevalence of the polymorphisms of this gene is consistent with global data for the respective races.

In clinical practice, pharmacogenetic testing of the *CYP2C19* gene is used to reduce adverse drug events in patients with cardiovascular diseases. Mirzaev *et al.* estimated the prevalence of alleles in Russian patients with acute coronary syndrome (ACS) in the Moscow region and Siberia. Statistically significant differences were found in *CYP2C19*3* allele frequency between the Russian ethnic group patients from eastern and central Siberia (p = 0.001; odds ratio = 1.05 [95% CI: 1.01–1.09]) [39]. Barysheva *et al.* used the results of pharmacogenetic testing of *CYP2C19* gene to personalize antiplatelet therapy in patients with ACS [19]. Fedorinov *et al.* studied the prevalence of alleles of this gene in Russian and Yakut patients with ACS and peptic ulcer disease in order to personalize therapy [45].

Muslimova *et al.* found association between carriage of *CYP2C19* 681A allele and a higher degree of platelet aggregation induced by Adenosine diphosphate in patients with coronary heart disease [46]. Golukhova *et al.* studied

Region of the Russian Federation Ethnic group Number of patients Dagestan Republic Laks 46 Healthy Dagestan Republic Laks 50 Healthy Dagestan Republic Dargins 50 6 Healthy European part Moscow Russians 59 6astric or du Furopean part Moscow Russians 59 6astric or du of Russia Russians 352 Healthy of Russia Russians 83 B-cell chroni in Russians Russians 83 B-cell chroni in Russians Russians 81 ACS in Russians 81 ACS 177	Characteristics of the population Healthy Gastric or duodenal peptic ulcer	Wt/Mt	\$2		∢	Allelic variants	ints				Ref.
Patients Laks 46 Laks 46 Avars 90 Dargins 50 Moscow Russians 971 Russians 971 76 Russians 83 177 Russians 83 177 Russians 83 177 Russians 81 76 General population 21 General population 55 Kursk 66 General population 561	lithy tric or duodenal peptic ulcer	Wt/Mt	*2								
Laks 46 Avars 46 Avars 90 Dargins 50 Dargins 50 Moscow Russians 971 Russians 971 76 Russians 83 76 Russians 83 76 Russians 83 76 Russians 81 76 General population 21 76 Kursk General population 55 Kursk General population 561	lithy tric or duodenal peptic ulcer	Wt/Mt	I			÷			*17		
Laks 46 Avars 90 Avars 90 Avars 90 Dargins 50 Moscow Russians 59 Moscom Russians 352 Russians 352 Russians 352 Russians 352 Russians 83 Russians 83 Russians 83 Russians 81 General population 21 General population 55 Kursk General population Starts 51	lithy tric or duodenal peptic ulcer		Mt/Mt	%	Wt/Mt	Mt/Mt	%	Wt/Mt	Mt/Mt	%	
Avars 90 Dargins 50 Dargins 50 Moscow Russians 59 Russians 971 76 Russians 83 76 Russians 81 76 Russians 71 71 Russians 81 76 Russian	tric or duodenal peptic ulcer	11	1	14.13	2	0	2.17	17	1	20.65	[5]
Dargins 50 Moscow Russians 59 Russians 971 Russians 352 Russians 352 Russians 81 Total 76 Russians 81	tric or duodenal peptic ulcer	20	-	12.22	0	0	0.00	30	m	20.00	
Moscow Russians 59 Russians 971 Russians 352 Russians 352 Russians 83 Russians 81 Russians 81 General population 21 Kutsk General population Kutsk General population	tric or duodenal peptic ulcer	5	0	5.00	0	0	0.00	22	-	24.00	
Russians971Russians352Russians352Russians83767617776Russians81General population21General population21General population55General population55General population51		13	2	14.40	-	0	0.84	19	-	17.79	[37]
Russians352Russians83Russians76776Russians81General population24General population21General population124General population55General population51	Gastric or duodenal peptic ulcer	245	14	14.00	10	1	0.62	399	66	27.34	[38]
Russians8376778181Russians81General population21General population124General population6eneral population6eneral population6eneral population6eneral population66eneral population<	lthy	84	4	13.10	I	I	I	I	I	I	[2]
76 177 Russians 81 General population 94 General population 124 General population 55 General population 561	B-cell chronic lymphocytic leukemia	18	0	10.84	I	I	I	I	I	I	[8]
177 Russians 81 General population 94 General population 21 General population 55 General population 51	Non-Hodgkin lymphoma	12	0	7.89	I	I	I	I	I	I	
Russians 81 General population 94 General population 21 General population 55 General population 55 General population 561	lthy	33	5	12.14	I	I	I	I	I	I	
General population 94 General population 21 General population 124 General population 55 General population 561		13	0	8.00	0	0	0.00	23	-	15.40	[39]
General population 21 General population 124 General population 55 General population 561	Pre-PCI patients	24	e	15.95	I	I	I	I	I	I	[40]
General population 124 General population 55 General population 561	Alcohol withdrawal	3	0	7.14	-	0	2.38	6	1	26.19	[41]
General population 55 General population 561		43	0	17.33	I	I	I	I	I	I	[42]
General population 561	ole CHD	14	2	16.36	I	I	I	I	I	I	[43]
	0	106	7	10.70	I	I	I	I	I	I	[25]
694 Healthy	lithy	130	21	12.40	Т	I	I	I	I	I	
General population 425 Essential hyp	Essential hypertension	80	4	10.40	I	I	I	I	I	I	[26]
391 Healthy	lithy	74	11	12.30	I	I	I	I	I	I	
Voronezh Russians 290 Healthy	llthy	56	5	11.37	2	0	0.34	I	T	I	[10]
Kazan Tatars 130 Healthy	lithy	n.d.	n.d.	11.90	I	I	I	n.d.	n.d.	21.20	[44]
Karachay-Cherkessia Circassians 77 Healthy	lithy	19	5	18.80	I	I	I	I	I	I	[13]
Karachays 125		31	2	14.00	I	I	I	I	I	I	
Siberia Surgut Russians 87 ACS		22	2	14.90	-	0	0.50	28	15	33.33	[39]
Novosibirsk, Kemerovo 222		39	0	11.50	0	0	0.00	62	7	17.10	
Irkutsk 122		14	1	10.60	5	-	2.80	43	9	22.20	

Region of th	Region of the Russian Federation	Ethnic group	Number of	Characteristics of the population					Allelic variants	nts				Ref.
			patients			*2			e. *			*17		
					Wt/Mt	Mt/Mt	%	Wt/Mt	Mt/Mt	%	Wt/Mt	Mt/Mt	%	
	Yakutsk	Russians	143	ACS	22	-	8.39	10	0	3.50	I	I	I	[45]
			63	Gastric or duodenal peptic ulcer	17	0	13.49	10	0	7.94	I	I	I	
		Yakuts	268	ACS	06	2	17.53	21	0	3.92	I	I	I	
			141	Gastric or duodenal peptic ulcer	41	0	14.54	22	0	7.80	I	I	I	
		Yakuts	229	Healthy	59	12	18.12	14	0	3.05	I	I	I	[15]
	Cheriktei	Yakuts	88	Healthy	27	7	23.30	8	0	4.60	I	I	I	[16]
	Kyzil	Tuvinians	88		24	-	14.80	4	0	2.30	I	I	I	
	Buriatia	Buryats	88		31	m	21.00	10	1	6.80	I	I	I	
	Altai	Altaians	87		20	e	14.90	7	0	4.00	I	I	I	
	Tomsk	Russians	87		16	2	12.20	0	0	0.00	I	I	I	
		General population 249	249	Breast cancer	109	4	23.49	5	0	1.01	I	I	I	[17]
		General population	166	IHD	39	e	13.55	I	I	I	I	I	I	[46]
Far East	Khabarovsk	Nanai	70	Healthy	28	m	24.30	12	0	8.60	m	0	2.10	[18]
Ural	Chelyabinsk	General population	63	Cardiac patients	19	2	11.50	I	I	I	41	m	23.50	[19]

residual platelet activity in patients receiving dual antiplatelet therapy. The frequency of the *CYP2C19*2* allele carriage in the high residual platelet reactivity group was significantly higher when compared with the group with normal platelet reactivity (40.0 vs 20.3%; p = 0.0035) [40]. Golukhova *et al.* studied sensitivity to clopidogrel in patients undergoing percutaneous coronary intervention (PCI). Platelet reactivity is higher in patients with heterozygous and homozygous carriers of *CYP2C19*2* versus common genotype [43]. Bokeria *et al.* used pharmacogenetic testing for the selection of antiplatelet therapy in patients with coronary artery disease [42]. Polonikov *et al.* found that *CYP2C19* rs4244285 GG was associated with an increased risk of essential hypertension (OR: 3.34; 95% CI: 1.48–7.51; p = 0.004) [26]. The same authors have tried to evaluate the association between the *CYP2C19* gene polymorphism and risk of coronary heart disease in Russian population, but the results were negative [25].

Pharmacogenetic testing is also used to improve the effectiveness of therapy in patients with diseases of the gastrointestinal tract. Sychev *et al.* found that the high frequency of *CYP2C19* genotypes associated with modified response to proton pump inhibitors in Russian patients with peptic ulcers [38]. Denisenko *et al.* found a connection between particular *CYP2C19* genotypes and urine metabolic ratio of Omeprazole in Russian peptic ulcer patients [37].

Zastrozhin *et al.* studied the relationship between carriage of alleles of various allelic variants of the *CYP2C19* gene and the development of adverse drug events in patients with alcohol withdrawal syndrome, taking Bromdihydrochlorphenylbenzodiazepine (Phenazepam[®]) [41].

Seredina *et al.* applied pharmacogenetic testing to increase the effectiveness of neoadjuvant chemotherapy in patients with breast cancer [17], and Gra *et al.* – for a personalized approach to therapy for patients with chronic lymphocytic leukemia and non-Hodgkin's lymphoma [8].

CYP2D6

From the standpoint of pharmacogenetics, the cytochrome CYP2D6 enzyme is a rather complex object of study due to the high polymorphism of its gene. The number of studied allelic variants reached 113, but not all of them are significantly associated with a decrease in the function of the enzyme [47]. In addition, this gene is characterized by an increase in the number of its copies, which increases the expression level of the enzyme [48]. Another feature of this enzyme is quite a wide range of changes in the activity. Statins, being a substrate of another isoenzyme, can reduce activity of CYP2D6 cytochrome that has been demonstrated *in vitro* study [49]. But what is more significant, the substrates of this isoenzyme can change its activity in the process of its metabolism, therefore, the effectiveness of the drug may suddenly decrease after some time from the start of treatment [50]. These subtle patterns of pharmacokinetics must be taken into account, since this cytochrome metabolizes 25% of drugs used nowadays [51].

Currently the *4 allele of this gene is the most studied is due to its high prevalence among representatives of the European race – 18.17%. The prevalence frequency among other populations: American – 10.76%, south/central Asian – 7.87%, Middle Eastern – 7.8%, African–American – 6.39%, African – 3.34%, Oceanian – 2.48% and east Asian 0.65% [52].

The analysis of the work of the Russian researchers showed a high prevalence of the allelic variant $CYP2D6^{*4}$ both in the general population and in individual ethnic groups. The highest prevalence was observed in Russians (17–20%), the lowest in the Mongoloid race (Nanai – 1.7%).

Cumulative information about the main characteristics of the studied population and the allele frequencies of the studied gene are presented in Tables 4 and 5.

Due to the broad class of drugs metabolized by CYP2D6 pharmacogenetic testing finds application in various diseases.

Ivanova *et al.* have reported an association between carriage of polymorphic variant 1846G>A *CY2D6*4* and genotype A/A of *CYP2D6* gene (responsible for debrisoquin-4-hydroxylase synthesis) with limbotruncal tardive dyskinesia in patients with schizophrenia receiving antipsychotics for a long time [62]. Sychev *et al.* found a statistically significant association between extrapyramidal disorders development and heterozygous *1846GA* genotype and *1846A* allele carrier frequency among all schizophrenic patients and among those of Tatars [60]. Zastrozhin *et al.* discovered that the lower efficacy and safety of Fluvoxamine in patients with depressive disorder and comorbid alcohol use disorders with *GA* genotype in *CYP2D6 1846G>A* polymorphic marker [55] and used test results to personalize therapy in patients with alcohol withdrawal syndrome [41]. Sychev *et al.* showed the correlations between the activity of CYP2D6 isozyme and the efficacy and safety of Haloperidol in patients with alcohol addiction [53,56].

Region of the RussianHunde of population patientCharacteristic of the patientDegestan RepublicLaks46HealthyDegestan RepublicLaks46HealthyDerotean part of RussianSoSoAccordinationDerotean part of RussianSoSoHealthyUnocowRusiansSoNon-Hodgkin lymphoonsUnocowRusiansSoHealthyUnocowRusiansSoHealthyUnocowRusiansSoHealthyUnocowGeneral populationSoHealthyUnocowGeneral populationSoAction lymphoonsUnocowGeneral populationSoAction lymphoonsUnocowRussiansLuocowSoAction lymphoonsUnocow <th></th> <th></th> <th></th> <th>Δ</th> <th>Allelic variants</th> <th></th> <th></th> <th></th> <th></th>				Δ	Allelic variants				
n Republic Laks 46 Avars 90 90 Avars 20 90 Avars 20 50 Dargins 50 50 w Russians 352 w General population 177 General population 45 173 General population 45 1230 Vasians Russians 1230 Russians Russians 1230 Petersburg Russians 1230 Russians Russians 1230 Petersburg Russians 1230 Russians Russians 1230 Patars Russians 1230 Russians Russians 1230 Russians Russians 1230 Russians Russians 1230 Russians Russians 124 Russians Russians 124 Russians Russians 124 Russians Russians 124 Russians Russians				ţ					Ref.
n Republic Laks 46 Avars 86 Avars 80 Avars 80 In part of Russians 352 w Russians 37 Ceneral population 81 General population 81 Status 82 Russians 130 Petersburg Russians Russians 130 Patars 82 Russians 37 Patars 216 Russians 37 Patars 216 Russians 37 Patars 216 Russians 216 Pay-Cherkessia 215 Russians 215 Pay-Cherkessia 216 Russians 215 Pay-Cherkessia 216 Russians			*3		*4			*10	
n RepublicLaks46Aars9050In part of Russians5050n part of Russians83352wRussians352wRussians352wRussians37Part of Russians7070Russians7070General population71General population81General population81General population81General population81Russians130Russians130PetersburgRussiansRussians131PetersburgRussiansRussians131PetersburgRussiansRussians131PetersburgRussiansRussians131PetersburgRussiansRussians131PetersburgRussiansRussians131PetersburgRussiansRussians131PetersburgRussiansRussians131PetersburgRussiansRussians131PetersburgRussiansRussians132PetersburgRussiansRussians132PetersburgRussiansPetersburg215PetersburgRussiansPetersburgRussiansPetersburgRussiansPetersburgRussiansPetersburgRussiansPetersburgRussians </th <th></th> <th>Wt/Mt Mt/Mt</th> <th>'Mt %</th> <th>Wt/Mt</th> <th>Mt/Mt</th> <th>W %</th> <th>Wt/Mt M1</th> <th>Mt/Mt %</th> <th></th>		Wt/Mt Mt/Mt	'Mt %	Wt/Mt	Mt/Mt	W %	Wt/Mt M1	Mt/Mt %	
Avars 90 Dargins 50 In part of Russians 50 w Russians 352 w Russians 37 General population 70 General population 81 Varians 1230 w Russians 1230 w Russians 1230 w Russians 131	Healthy	1	I	17	0	18.47	I	1	[5]
Dargins 50 In part of Russians 352 w Russians 352 w Russians 352 Russians 8 352 Russians 8 352 Russians 8 352 Russians 76 76 Russians 76 75 General population 70 76 General population 70 76 General population 70 75 Vetersburg Russians 71			ı	23	0	12.77 _	T	T	
nn part of Russians 352 w Russians 352 w Russians 352 Russians 352 352 Russians 352 352 Russians 352 352 Russians 352 352 Russians 36 37 Russians 31 37 Russians 37 37 Vetersburg Russians 37 Russians 37 37				19	0	19.00 _		I	
wRussians352Russians83Russians83Russians76Antil Russians177General population70General population81General population81General population81General population81General population81General population81General population81General population81General population81General population81Russians123Patars82Russians37Patars82Russians37Patars82Russians37Patars82Russians77Russians77Russians77Russians77Russians77Russians77Russians77Russians77Russians77Russians77Russians77Russians77St<									
Russians 8 Russians 76 Reneral population 70 General population 70 General population 81 Russians 1230 Varsians 181 Russians 181 Russians 181 Paters Russians Russians 171 Paters 181 Russians 173 Paters 181 Paters 173 Russians 173 Paters 174 Paters 174 <td></td> <td>6 0</td> <td>06.0</td> <td>91</td> <td>14</td> <td>17.00 -</td> <td>1</td> <td>·</td> <td>[7]</td>		6 0	06.0	91	14	17.00 -	1	·	[7]
76 177 6eneral population 70 6eneral population 71 8 8 8 8 8 8 9 </td <td>B-cell chronic lymphocytic leukemia</td> <td>1</td> <td>0.60</td> <td>24</td> <td>m</td> <td>18.07 -</td> <td>1</td> <td>I</td> <td>[8]</td>	B-cell chronic lymphocytic leukemia	1	0.60	24	m	18.07 -	1	I	[8]
177 General population 70 General population 81 General population 21 Kussians 1230 Russians 181 Russians 171 Russians 290 Russians 214 Russians 214 Russians 214 Russians 214 Russians 215 Cherkessia Circassians Russians 175 Sav-Cherkessia Circassians Save and Chita General population Save and Chita General population <td></td> <td>1 0</td> <td>0.65</td> <td>20</td> <td>m</td> <td>17.10 -</td> <td>T</td> <td>ı</td> <td></td>		1 0	0.65	20	m	17.10 -	T	ı	
General population 70 General population 81 General population 45 General population 45 General population 69 General population 71 Varsians 1230 Russians 17 Pathon Russians Russians 21 Pathon Russians Pathon 17 Pathon 200 Pathon 21 Pathon 215 Pathon 215 Pathon 215 Pathon 215 Pathon 21		4 1	1.69	49	∞	18.36 _	T	ı	
General population 81 General population 45 General population 69 General population 69 General population 69 General population 69 Russians 1230 Russians 181 Russians 17 Russians 214 Russians 215 Russians 17 Russians 175	Alcohol addiction		ı	n.d.	.p.u	12.14 _	T	ı	[53]
General population 45 General population 69 General population 69 General population 21 Russians 1230 Russians 181 Russians 290 N Russians Russians 214 Russians 214 Russians 215 Russians 215 Russians 177 Russians 175 Russians 176 Russians 178 Russians 178 Russians 177 Russians 178 Russians 178 Russians 178 Russians 178 Russians 178	Arterial hypertension		ı	I		74	1 7	54.32	[54]
General population 69 etersburg General population 21 etersburg Russians 1230 Russians Russians 181 Russians Russians 181 Russians Russians 181 Russians Russians 181 Russians Russians 17 Russians Russians 290 Russians Russians 215 Russians Ceneral population 214 Russians Circassians 77 Russians Circassians 77 Russians Circassians 125 Russians General population 214 Russians Circassians 77 Russia	Depressive disorder and comorbid alcohol use disorder	1	I	17	0	18.89	I	ı	[55]
General population21etersburgRussians1230etersburgRussians1230RussiansRussians17ezhRussians290nRussians290nRussians27etersburgRussians215nTatars214nGeneral population214ay-CherkessiaCircassians77ay-CherkessiaCircassians77ovo and ChitaGeneral population215skRussians77sibirskRussians77sibirskIndra Nentsi108	Alcohol use disorder		1	17	0	12.32 _	1	I	[56]
vetersburgRussians1230RussiansRussians181RussiansGeneral population17Russians29037Russians3737Russians3737Russians3737Russians3737Russians3737Russians3737Russians214214Russians77214	Alcohol withdrawal			4	0	9.52 -	1	I	[41]
Russians 181 Russians 17 Beneral population 17 Beneral population 17 Prastians 290 Russians 37 Russians 37 Patars 290 Russians 37 Russians 216 General population 215 Addition 214 Russians 77 Starstow 214 Nay-Cherkessia 6rcassians Cherkessia 77 Starstow 125 Starstow 125 Starstow 93 Starstow 93 Starstow 108 Starstow 108 Starstow 108		n.d. n.d	1.30	n.d	, b.n	17.70 _	T	T	[57]
General population 17 azh General population 290 n Russians 290 n Russians 37 Tatars 215 General population 215 ay-Cherkessia Circassians 77 ay-Cherkessia Circassians 77 ow and Chita General population 475 ow and Chita General population 475 sk Russian 93 ibirsk Tundra Nentsi 108		в 0	1.66	54	m	16.57 8	-	2.76	[58]
azhRussians290nRussians37Tatars837Tatars837Tatars6215General population215ay-Cherkessia77ay-Cherkessia77sy-Cherkessia77core and Chita6ovo and Chita6skRussianskRussiansk8ibirsk108ibirskTundra Nentsi102		0 8	8.82	5	0	14.70 -	I	I	[59]
N Russians 37 Tatars 42 Tatars 42 General population 215 General population 214 Nay-Cherkessia 77 Nay-Cherkessia 77 Nay-Cherkessia 77 Nay-Cherkessia 77 State 8 Nay-Cherkessia 77 State 77 Nay-Cherkessia 77 State 77 Nay-Cherkessia 77 State 8 State 93 State 70 State 70 State 70 State 70 State 70 State 71		6 0	1.03	87	6	18.10 22	-	4.13	[10]
Tatars 42 General population 215 General population 214 nay-Cherkessia 77 ay-Cherkessia 77 Rarachays 175 ovo and Chita General population sk Russian Yakuts 93 tibrsk Tundra Nentsi tibrsk Tundra Nentsi	Schizophrenia		ī	12	e	24.30	T	T	[09]
General population 215 iay-Cherkessia Circassians 77 iay-Cherkessia Trachays 77 iay-Cherkessia Circassians 77 owo and Chita General population 475 owo and Chita General population 93 sk Russian 93 ibirsk Tundra Nentsi 108		1	I	10	0	11.90 _	T	I	
214 nay-Cherkessia Circassians 77 Nay-Cherkessia Transmit of the strain	Asthma			64	. L	18.14 -	1		[61]
iay-Cherkessia Circassians 77 Karachays 125 Karachays 125 ovo and Chita General population 475 sk Russian 93 Yakuts 108 tibirsk Tundra Nentsi 102	Healthy	1	I	62	9	17.29 –	I	I	
Karachays 125 ovo and Chita General population 475 sk Russian 93 Yakuts 108 tibirsk Tundra Nentsi 102	Healthy		,	19	7	15.00 -	1	I	[13]
ovo and Chita General population 475 sk Russian 93 Yakuts 108 ibirsk Tundra Nentsi 102			ı	38	7	20.80 -	T	ı	
General population 475 Russian 93 Yakuts 108 Tundra Nentsi 102									
Russian 93 Yakuts 108 Irsk Tundra Nentsi 102		8 1	1.05	86	11	11.36 -	T	T	[62]
Yakuts 108 Tundra Nentsi 102	CHD			20	0	10.75 30	0	16.13	[63]
Tundra Nentsi 102		ı ı	ı	22	0	10.19 35	0	16.20	
	Healthy		ı	p.u	p.u	7.00 -	I	·	[64]
General population 96				n.d	p.u	20.00 -	1	ı	
Far East:									
– Khabarovsk Nanai 70 Healthy	Healthy		1	2	0	1.40 _	1	ı	[18]

Region of the Russian	Ethnic group	Number of	Characteristics of the population		Allelic varia	nts	Ref.
Federation		patients		Wt/Mt	Mt/Mt	%	
CYP2D6*2							
Saint Petersburg	Russian	1230	Healthy	n.d.	n.d	35.7	[57]
CYP2D6*5							
Voronezh	Russian	290	Healthy	15	0	2.58	[10]
Saint Petersburg		1230		n.d.	n.d.	1.60	[57]
CYP2D6*6							
Voronezh	Russian	290	Healthy	7	0	1.20	[10]
Saint Petersburg		1230		n.d.	n.d.	1.00	[57]
CYP2D6*7							
Saint Petersburg	Russian	1230	Healthy	n.d.	n.d.	0.04	[57]
CYP2D6*41							
Saint Petersburg	Russian	1230	Healthy	n.d.	n.d.	7.60	[57]

Often pharmacogenetic testing is necessary with the simultaneous prescribing drugs from different classes. Goryachkina *et al.* first showed that metabolism of metoprolol in patients with acute myocardial infarction (AMI) can be inhibited by the simultaneous administration of paroxetine for the treatment of depressive disorder [59]. However, even in the absence of paroxetine, metabolism of metoprolol in patients with AMI can be accelerated as a result of doubling of the *CYP2D6* gene and increased expression of the enzyme [58]. In patients with arterial hypertension, the carriage of certain alleles of this gene can increase the effectiveness of therapy with betaxolol [54].

Polonikov *et al.* first rated gene–gene and gene–environment interactions for disease susceptibility for asthma [61]. Fedorinov *et al.* revealed an association between the polymorphic marker G1846A carrier and the lower effective dose of Bisoprolol in patients with CHD [63].

CYP2D6 gene polymorphisms have also been studied to create a biochip as a tool for predictive analysis in oncohematology [8].

CYP1A1

As for the cytochrome *CYP1A1* gene, 13 allelic variants have been discovered, and 12 SNPs the gene of this isoenzyme require further study [65]. The role of this isoenzyme in carcinogenesis processes is discussed, in view of its involvement in the metabolic activation of procarcinogens (polycyclic aromatic hydrocarbons and aromatic amines) into reactive metabolites. It also participates in the metabolism of steroidal hormones including estrogens [66]. In view of the little scrutiny, it is difficult to form a reliable idea of the prevalence rate of this gene in the world population.

Cumulative information about the main characteristics of the studied population and the frequency of *CYP1A1* alleles are presented in Table 6.

As described above, most of the research is focused on oncology. Mikhailova *et al.* were the first to study this problem in Russia. The authors found no statistically significant differences in the prevalence frequency of the allelic variant *CYP1A1*2A* in patients with benign and malignant tumors compared with healthy volunteers [70].

Belogubova *et al.* have shown the association of allelic variant *CYP1A1*2* with a predisposition to the development of squamous cell lung carcinoma [69]. Gra *et al.* found that the allelic variant of *CYP1A1*2B* was more common in patients with B-cell chronic lymphocytic leukemia [8].

Pulmonology was another area that was covered in the Russian studies of CYP1A1. Polonikov *et al.* found that an absence of genotypes 462IV of the *CYP1A1* gene (p = 0.050) and also possessing the GSTM1-null genotype (p = 0.023) were significantly associated with an increased risk of asthma only in smokers [61]. Salnikova *et al.* proved that allelic variants of *CYP1A1*2A* and *CYP1A1*2B* increase the patient's sensitivity to the community-acquired and nosocomial pneumonia [68] disease and increase the risk of complications of these diseases [67].

Table 6.	Allele and	Table 6. Allele and genotype frequencies of	quencies of CYP1A1 gene in different ethnic groups in Russia.	lene in (differen	t ethnic	groups	in Russ	ia.							
Region of	Ethnic group		<u>.</u>						Allelic variants	ariants						Ref.
the Russian Federation		patients	population		*2A			*2B			*2C			*4		
				Wt/Mt	Mt/Mt	%	Wt/Mt	Mt/Mt	%	Wt/Mt	Mt/Mt	%	Wt/Mt	Mt/Mt	%	
European pa	European part of Russia															
Moscow	Russians	352	Healthy	34	m	5.68	36	-	5.40	I	I	I	13	0	1.90	[7]
	Russians	83	B-cell chronic lymphocytic leukemia	11	0	6.62	12	0	7.89	I	I	I	7	0	4.21	[8]
		76	Non-Hodgkin lymphoma	10	0	6.57	10	0	6.57	I	I	I	4	0	2.63	
		177	Healthy	18	1	5.64	14	0	4.05	I	I	I	6	1	3.10	
	General population	341	Community-acquired pneumonia	59	+	8.94	I	I	I	33	0	4.84	I	I	I	[67]
	General population	334	Community-acquired and nosocomial pneumonia	57	+	8.83	I	I	I	32	0	4.79	I	I	I	[68]
		141	Healthy	33	0	12.31	I	I	I	8	0	2.99	I	I	I	
Kursk	General population	215	Asthma	40	1	9.77	I	I	I	33	-	8.14	I	I	I	[61]
		214	Healthy	51	1	12.38	I	I	I	30	1	7.48	I	I	1	
Saint Petersburg	General population	246	Elderly donors	49	1	10.37	I	I	I	I	I	I	I	I	I	[69]
		204	Middle-aged donors	41	2	11.03	I	I	I	I	I	I	I	I	I	
		141	Lung cancer	35	2	13.83	I	I	I	I	I	I	I	I	I	
Ufa	Russians	319	Healthy	66	e	12.63	I	I	I	28	1	4.78	I	I	I	[12]
	Tatars	279		60	8	16.38	I	I	I	27	2	6.38	I	I	I	
	Bashkirs	144		52	9	23.53	I	I	I	28	0	10.45	I	I	I	
Siberia																
Novosibirsk	General population	166	Malignant tumor	36	0	10.71	I	I	I	I	I	I	I	I	I	[0/]
		170	Benign tumor	28	0	8.24	I	I	I	I	I	I	I	I	I	
		170	Healthy	40	0	11.90	I	T	I	I	I	Т	I	I	T	

Region of the Russian	Ethnic group	Number of	Characteristics of the population		Allelic varia	nts	Ref
Federation		patients		Wt/Mt	Mt/Mt	%	
CYP2C8*1B rs7909236							
Kursk	Common population	561	CHD	210	35	25.00	[25
		694	Healthy	239	33	22.00	
		425	Essential hypertension	151	32	25.30	[26
		391	Healthy	118	11	17.90	
СҮР2С8*2 Т805А							
Tomsk	Common population	244	Breast cancer	1	0	0.20	[17]
CYP2C8*3 G416A + A1196G	i						
Tomsk	Common population	247	Breast cancer	44	0	8.90	[17
CYP2C8 rs1934953							
Kursk	Common population	561	CHD	255	61	36.60	[25
		694	Healthy	314	64	31.80	
		425	Essential hypertension	197	50	34.90	[26
		391	Healthy	181	29	30.60	

CYP2C8

One of the poorly studied isoenzymes is cytochrome CYP2C8. Currently, 14 allelic variants of the gene of this isoenzyme have been discovered [71]. This isoenzyme is involved in the biotransformation of drugs such as amiodarone, cerivastatin, paclitaxel, repaglinide, rosiglitazone, diclofenac, ibuprofen, fluvastatin, morphine, carbamazepine, verapamil – among others. [72]. The most common is the *CYP2C8*2* allele, with its prevalence in black populations reaching 18% and about 1% in white subjects [73].

Cumulative information about the main characteristics of the studied population and the frequency of *CYP2C8* alleles are presented in Table 7.

The allelic variants of *CYP2C8*1B* rs7909236 and *CYP2C8* rs1934953 were studied in two papers by Russian researchers. Polonikov *et al.* studied these polymorphisms from the point of view of the risk of developing coronary heart disease in the Russian population [25] and established their relationship with the risk of developing essential arterial hypertension [26].

Seredina *et al.* studied the *CYP2C8*2* T805A and *CYP2C8*3* G416A + A1196G polymorphisms from the standpoint of the effectiveness of neoadjuvant chemotherapy in patients with breast cancer, but no reliable association data was obtained [17].

Other oxidation enzymes

Besides CYP450, other enzymes are also involved in drug metabolism.

For example, the human CES1, also localized in the liver, is involved in the metabolism of dabigatran [74]. Sychev *et al.* did not confirm the effect of the *CES1* rs2244613 polymorphism Dabigatran equilibrium peak concentration in patients after total knee arthroplasty [75].

Another enzyme is SULT1A1, which is involved in estrogen metabolism [76]. Mikhailova *et al.* proved the association between the carriage of the G allele of *SULT1A1*2* with hormone-dependent cancers in women [70].

Cumulative information about the main characteristics of the studied population and the frequency of *CES1* and *SULT1A1* alleles are presented in Table 8.

Membrane transporters

In addition to the CYP450 system, membrane transporters play an important role in drug metabolism. With the help of them, the drug is transported into the cell and the drug is excreted into the extracellular space.

Cumulative information about the main characteristics of the studied population and the frequency of *SLCO1B1* and *ABCB1* alleles are presented in Tables 9 and 10.

enotype frequencie	s of CES1 and	SULT1A1 genes in different e	ethnic gro	oups in R	lussia.	
Ethnic group	Number of	Characteristics of the population		Allelic varia	nts	Ref.
	patients		Wt/Mt	Mt/Mt	%	
General population	60	Total knee arthroplasty	21	6	27.5	[75]
General population	166	Healthy	70	28	37.50	[70]
	170	Malignant tumor	86	16	34.71	
	170	Benign tumor	60	50	47.06	
	Ethnic group General population	Ethnic group Number of patients General population 60 General population 166 170	Ethnic group Number of patients Characteristics of the population General population 60 Total knee arthroplasty General population 166 Healthy 170 Malignant tumor	Ethnic group Number of patients Characteristics of the population Wt/Mt General population 60 Total knee arthroplasty 21 General population 60 Healthy 70 Image: The second sec	Ethnic group Number of patients Characteristics of the population with Mt/Mt Allelic varia General population 60 Total knee arthroplasty 21 6 General population 60 Total knee arthroplasty 21 6 General population 166 Healthy 70 28 170 Malignant tumor 86 16	Wt/MtMt/Mt%General population60Total knee arthroplasty21627.5General population166Healthy702837.50170Malignant tumor861634.71

Region of the	Ethnic group	Number of	Characteristics of the population			Allelic	variants			Ref.
Russian Federation		patients		SLC	01B1 +52	1T>C	Al	BCB1 34350	T<2	
				Wt/Mt	Mt/Mt	%	Wt/Mt	Mt/Mt	%	
Dagestan Republic	Laks	46	Healthy	7	1	9.78	21	21	67.47	[5]
	Avars	90		26	0	14.44	48	26	55.55	
	Dargins	50		8	1	10.00	21	21	63.00	
European part of Rus	sia:									
– Moscow	Russians	50	Atrial fibrillation	-	-	-	25	15	55.00	[77]
	General population	17	Atrial fibrillation and acute stroke	-	-	-	9	3	44.12	[78]
	General population	341	Community-acquired pneumonia	-	-	-	141	66	40.38	[67
	General population	334	Community-acquired and nosocomial pneumonia	-	-	-	140	53	36.83	[68
		141	Healthy	-	-	-	63	31	46.64	
		21	Alcohol withdrawal	-	-	_	9	1	16.19	[41]
		100	Stage I–II hypertension (HT)	-	-	-	53	30	56.50	[79
	Russians	1071	Hypercholesterolemia	346	60	21.76	_	_	-	[80]
	Yakuts	76		11	3	11.18	-	-	-	
	General population	50	Valvular atrial fibrillation	_	_	_	25	15	55.00	[81]
– Yaroslavl	General population	81	Statin taking	27	2	19.14	_	_	-	[82]
– Ryazan	General population	121	Statin taking	46	8	26.70	-	_	-	[83]
– Voronezh	General population	285	Colorectal cancer	-	-	-	129	70	47.20	[84
		275	Non cancer(not healthy)	-	-	-	140	78	53.80	
– Saratov	General population	60	Total knee arthroplasty	_	-	_	29	16	50.8	[75]
Siberia:										
– Yakutsk	Yakuts	100	Healthy	12	1	14.00	-	_	-	[15]
Far East:										
– Khabarovsk	Nanai	70	Healthy	25	4	23.60	39	12	45.00	[18

ABCB1 (MDR1, P-gp)

P-gp is a protein from the ABC family of transporters encoded by the *ABCB1* gene. This transporter is located on the membrane of enterocytes, regulating the process of absorption and release of the drug, as well as on the membrane of other cell types, for example, cells that form the blood-brain barrier [85]. The gene of this transporter has many different allelic variants, including those affecting the function of the transporter. To date, 1279 SNPs in the *ABCB1* gene region are open, 62 of which are coding [86].

In clinical studies, Sychev *et al.* established that TT genotype of rs1045642 polymorphism of the *ABCB1* gene was associated with higher dabigatran equilibrium peak concentrations and the higher risk of bleeding than the presence of CC genotype (p < 0.008) [75] and indicate the potential of pharmacogenetic testing for rs1045642 SNP when prescribing amlodipine for the first time in Caucasian patients with stage I–II arterial HT [79]. Osswald *et al.* found that life-long nonsmokers with an age \geq 63 years who carried the 3435CC or 3435TT genotype developed more frequently colorectal cancer than others (OR = 3.9; 95% CI: 2.0–7.7) [84]. Rozhkov *et al.* revealed

Region of the Russian	Ethnic group	Number of	Characteristics of the population		Allelic varia	nts	Ref.
Federation		patients		Wt/Mt	Mt/Mt	%	
ABCB1 rs4148738							
Moscow	General population	17	Atrial fibrillation and acute stroke	7	6	55.88	[78]
Saratov	General population	60	Total knee arthroplasty	25	15	45.8	[75]
ABCB1 -129T>C (rs3213619)							
Voronezh	General population	285	Colorectal cancer	16	0	2.80	[84]
		275	Non cancer(not healthy)	15	0	2.70	

that *ABCB1* CT and TT genotypes significantly associated with higher risk of bleeding in patients with valvular atrial fibrillation taking acenocoumarol [81]. Similar results were obtained in another study [77]. Zastrozhin *et al.* took into account the polymorphism of the gene *ABCB1* to create pharmacogenetic-guided personalization of drug dose in patients with alcohol withdrawal syndrome, due to its association with the development of adverse drug events [41]. Kryukov *et al.* found no effect of polymorphism on the pharmacokinetics of apixaban [78]. Salnikova *et al.* did not find an association of *ABCB1* gene polymorphisms with a predisposition to community-acquired and nosocomial pneumonia [68] and an increased incidence of complications [67].

Of the rarer alleles, only ABCB1 rs4148738 [32,44] and ABCB1 -129T>C (rs3213619) [84] were studied.

SLCO1B1

SLCO1B1 is a polypeptide that transports organic anions across a cell membrane. It acquired its significance in particular due to the participation in the metabolism of statins by transporting drugs into hepatocytes [87]. Currently, the definition of genotypes for the allelic variant *SLCO1B1*5* has already been recommended for practical use by experts from the European Science Foundation [88].

The polymorphisms of this gene remain virtually unexplored for the Russian population. There are data on the frequency of the prevalence of the allelic variant SLCO1B1 + 521T>C among Russians [80], Yakuts [15,80], Nanai [18], Laks, Avars and Darghins [5]. Khokhlov *et al.* studied the frequency of SLCO1B1 + 521T>C in patients taking statins. Adverse drug reactions developed in 19.5% when taking Simvastatin and carriage frequency of allele was 19.14% [82]. Solodun *et al.* found a statistically significant decrease in atherogenic cholesterol fractions in carriers of SLCO1B1*5 alleles while taking Atorvastatin [83].

Prospects for the use of pharmacogenetic testing in the Russian Federation

To date, the recommendations of professional communities regarding the use of pharmacogenetic testing are controversial and disappointing. A striking example is the recommendations of the American Heart Association and the American College of Cardiology, which contradict the position of the US FDA and the CPIC regarding the appropriateness of using *CYP2C19* pharmacogenetic testing for clopidogrel in patients with ACS [89]. Such inconsistencies are largely due to a lack of large comparative prospective clinical trials confirming the advantages of the pharmacogenetic approach to therapy as compared with the standard one, in other words, the expediency of using a marker in clinical practice.

According to the recommendations of the Evaluation of Genomic Applications in Practice and Prevention working group, clinical validity is the ability of a pharmacogenetic marker to predict the response to a drug, and clinical utility is the ability to improve clinical outcomes [90]. However, most pharmacogenetic markers overcome the barrier of clinical validity and only a few overcome the barrier of clinical utility. From these standpoints, the most clinically valid and clinically useful for therapeutic practice are gene/product pairs: CYP2C19 – clopidogrel, CYP2C9 and VKORC1 – warfarin, and SLCO1B1 – statins [91]. Pharmacogenetic testing for these markers has already been implemented in the standards of care at the University of Florida, where testing for the polymorphism of the *CYP2C19* gene is performed on all patients who have undergone cardiac catheterization since 2012, as well as at the University of Vanderbilt, where testing for the polymorphism of the *SLCO1B1* gene is performed prior to the prescription of statins [92].

Conclusion

The results of the systematic review presented in the current study indicate the importance of studying the most clinically valid and clinically useful pharmacogenetic markers (*CYP2C19, CYP2C9, VKORC1, SLCO1B1*) among different ethnic groups of the Russian Federation. For example, *CYP2C9*2* allelic variant prevails in Russians as representatives of the Europoid race, and is much rarer in Mongoloid race: Yakuts, Buryats, Tuvinians. Besides, the *CYP2C9*3* allelic variant is unusually widespread among representatives of ethnic groups of the Caucasus: Circassians, Laks, Avars and Dargins. The prevalence of *VKORC1* 1639G>A rs9923321 polymorphism in the Yakut population is two-times higher than among patients of the Moscow region – 83.14 and 36.26%, respectively. The analysis of the works of Russian researchers on the prevalence of *SLCO1B1+521T>C* also revealed important regional ethnic features.

With the accumulation of evidence of clinical validity and clinical utility of other pharmacogenetic markers (*CES1, CYP2D6*4*, etc.), the problem of interethnic differences in the presence of clinically significant polymorphisms of these genes, identified in earlier studies in the Russian Federation, requires increasing attention.

Executive summary

CYP450 enzymes

CYP2C9 + VKORC1 & GGCX

- Fifteen studies were conducted on allelic variants of CYP2C9*2, *3 of them: nine in the European part of Russia, three in Siberia, one in the Far East, the Urals and the Republic of Dagestan.
- In addition to healthy volunteers, associative studies were also conducted in cardiac and oncological patients.
- Minor allelic variants of CYP2C9 rs9332242, rs4918758, rs61886769 as well as other enzymes of warfarin biotransformation (VKORC1 and GGCX) are practically not studied in the Russian population.

CYP2C19

- The most studied at the moment are CYP2C19*2, *3, *17 allelic variants.
- Fourteen studies were conducted in the European part of Russia, six in Siberia, one in the Republic of Dagestan, the Far East and the Urals.
- Pharmacogenetic testing for this gene has found application in oncology, cardiology, psychiatry and gastroenterology.

CYP2D6

- The most significant allelic variants that are reflected in the studies are CYP2D6*3, *4, *10.
- Most of the investigations concentrated in the European part of Russia 14, 3 were held in Siberia, 11 in the republic of Dagestan and the Far East.
- Much research has focused on psychiatry and narcology. Cardiology, pulmonology and oncology are covered to a small extent.

CYP1A1

- Studies of the gene of this cytochrome were conducted among patients of the European part of Russia 7, and Siberia 1.
- Most research is focused on oncology.

CYP2C8

- One of the poorly studied isoenzymes.
- Researches in relation to cardiology and oncology were conducted on the general population.
- The prevalence frequency of allelic variants of this gene among certain ethnic groups of the Russian Federation remains unclear.

CES1 and SULT1A1

- One of the poorly studied enzymes.
- For these genes there is a total of one study in the general population of patients.
- It is necessary to conduct population-ethnic and associative clinical studies.

ABCB1 (MDR1, P-gp)

- The allelic variant ABCB1 3435C>T is studied in detail.
- However, the bulk of the research was conducted on the general population of patients in the European part of Russia.

SLCO1B1

- Was studied in just six studies.
- In addition to population-ethnic studies in healthy volunteers, attempts were made to personalize statin therapy based on pharmacogenetic testing.

In the Russian population, there are still no studies of such enzymes as uridine 5'-diphosphoglucuronosyltransferase, human flavin-containing monooxygenase, monoamine oxidases, aldehyde oxidase and such membrane transporters as SLCO1B3 and ABCG2.

These racial and ethnic peculiarities may affect the development and priority of the implementation of pharmacogenetic testing programs in different regions of the Russian Federation.

Supplementary data

To view the supplementary data that accompany this paper please visit the journal website at: https://www.futuremedicine.com/d oi/suppl/10.2217/pgs-2019-0013

Financial & competing interests disclosure

This study was supported by the Russian Science Foundation (Grant No. 18-75-00112) 'Genetic determinants of drug sensitivity to the rooted peoples of the North Caucasus, the Republic of Crimea, the Volga region and Siberia'. The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

No writing assistance was utilized in the production of this manuscript.

References

- 1. Hippocrates quotes. BrainyQuote (2019). www.brainyquote.com/authors/hippocrates
- Lu YF, Goldstein DB, Angrist M, Cavalleri G. Personalized medicine and human genetic diversity. *Cold Spring Harb. Perspect. Med.* 4(9), a008581 (2014).
- 3. Guidelines. Cpicpgx.org (2019). https://cpicpgx.org/guidelines/
- 4. All-Russian population census 2010. The national composition of the population of Russia (2010). www.gks.ru/free_doc/new_site/perepis2010/croc/perepis_itogi1612.htm
- 5. Mirzaev K, Sychev D, Ryzhikova K *et al.* Genetic polymorphisms of cytochrome P450 enzymes and transport proteins in a Russian population and three ethnic groups of Dagestan. *Genet. Test. Mol. Biomarkers* 21(12), 747–753 (2017).
- 6. Sychev D, Rozhkov A, Ananichuk A, Kazakov R. Evaluation of genotype-guided acenocoumarol dosing algorithms in Russian patients. *Drug Metab Pers Ther.* 32(2), 317–327 (2017).
- Gra O, Mityaeva O, Berdichevets I et al. Microarray-based detection of CYP1A1, CYP2C9, CYP2C19, CYP2D6, GSTT1, GSTM1, MTHFR, MTRR, NQO1, NAT2, HLA-DQA1, and AB0 allele frequencies in native Russians. Genet. Test. Mol. Biomarkers 14(3), 329–342 (2010).
- Gra O, Glotov A, Nikitin E *et al.* Polymorphisms in xenobiotic-metabolizing genes and the risk of chronic lymphocytic leukemia and non-Hodgkin's lymphoma in adult Russian patients. *Am. J. Hematol.* 83(4), 279–287 (2008).
- 9. Ivashchenko D, Rusin I, Sychev D, Grachev A. The frequency of *CYP2C9, VKORC1*, and *CYP4F2* polymorphisms in russian patients with high thrombotic risk. *Medicina* 49(12), 81–85 (2013).
- 10. Gaikovitch E, Cascorbi I, Mrozikiewicz P *et al.* Polymorphisms of drug-metabolizing enzymes CYP2C9, CYP2C19, CYP2D6, CYP1A1, NAT2 and of P-glycoprotein in a Russian population. *Eur. J. Clin. Pharmacol.* 59(4), 303–312 (2003).
- 11. Pchelina S, Sirotkina O, Taraskina A, Vavilova T, Shwarzman A, Schwartz E. The frequency of cytochrome *P450 2C9* genetic variants in the Russian population and their associations with individual sensitivity to warfarin therapy. *Thromb. Res.* 115(3), 199–203 (2005).
- 12. Korytina G, Kochetova O, Akhmadishina L, Viktorova E, Victorova T. Polymorphisms of cytochrome *P450* genes in three ethnic groups from Russia. *Balkan Med. J.* 8(2), 15–23 (2012).
- Romodanovsky D, Khapaev B, Ignatiev I, Kukes V, Karkischenko V. [Frequencies the «slow» allele variants of the genes coding isoenzymes of cytochrome *P450 CYP2D6, CYP2C19, CYP2C9* in Karachaevs and Circassians]. *Biomedicine* 2, 33–37 (2010) (In Russian).
- 14. Baturin V, Tsarukyan A, Kolodijchuk E. Study of *CYP2C9* gene polymorphism in ethnic groups of the population of stavropol region. *Med. News North Caucasus* 9(1), 51–58 (2014).
- 15. Vasilyev FF, Danilova DA, Kaimonov VS, Chertovskih YV, Maksimova NR. Frequency distribution of polymorphisms of *CYP2C19*, *CYP2C9*, *VKORC1* and *SLCO1B1* genes in the Yakut population. *Res. Pharm. Sci.* 11(3), 259–264 (2016).
- 16. Makeeva O, Stepanov V, Puzyrev V, Goldstein D, Grossman I. Global pharmacogenetics: genetic substructure of Eurasian populations and its effect on variants of drug-metabolizing enzymes. *Pharmacogenomics* 9(7), 847–868 (2008).
- 17. Seredina T, Goreva O, Talaban V, Grishanova A, Lyakhovich V. Association of cytochrome *P450* genetic polymorphisms with neoadjuvant chemotherapy efficacy in breast cancer patients. *BMC Med. Genet.* 13(1), 13–25 (2012).
- 18. Sychev D, Shuev G, Suleymanov S et al. Comparison of CYP2C9, CYP2C19, CYP2D6, ABCB1, and SLCO1B1 gene-polymorphism frequency in Russian and Nanai populations. *Pharmgenomics Pers. Med.* 10, 93–99 (2017).

- 19. Barysheva V, Ketova G. [Analysis of the prevalence of genotypes according to pharmacogenetic testing]. *Mod. Probl. Sci. Educ.* 4, 112–125 (2016) (In Russian).
- 20. PharmVar CYP2C9 (2019). www.pharmvar.org/gene/CYP2C9
- 21. Kirchheiner J, Brockmoller J. Clinical consequences of cytochrome P450 2C9 polymorphisms. *Clin. Pharmacol. Ther.* 77(1), 1–16 (2005).
- 22. WarfarinDosing. www.warfarindosing.org/Source/Home.aspx
- Johnson J, Caudle K, Gong L *et al.* Clinical Pharmacogenetics Implementation Consortium (CPIC) guideline for pharmacogenetics-guided warfarin dosing: 2017 update. *Clin. Pharmacol. Ther.* 102(3), 397–404 (2017).
- 24. PharmGKB. CYP2C9 frequency table. https://api.pharmgkb.org/v1/download/file/attachment/CYP2C9_frequency_table.xlsx
- 25. Polonikov A, Kharchenko A, Bykanova M *et al.* Polymorphisms of *CYP2C8*, *CYP2C9* and *CYP2C19* and risk of coronary heart disease in Russian population. *Gene* 627, 451–459 (2017).
- Polonikov A, Bykanova M, Ponomarenko I et al. The contribution of CYP2C gene subfamily involved in epoxygenase pathway of arachidonic acids metabolism to hypertension susceptibility in Russian population. Clin. Exp. Hypertens. 39(4), 306–311 (2017).
- 27. Cain D, Hutson S, Wallin R. Assembly of the warfarin-sensitive vitamin K 2,3-epoxide reductase enzyme complex in the endoplasmic reticulum membrane. *J. Biol. Chem.* 272(46), 29068–29075 (1997).
- Liang R, Li L, Li C et al. Impact of CYP2C9*3, VKORC1-1639, CYP4F2 rs2108622 genetic polymorphism and clinical factors on warfarin maintenance dose in Han-Chinese patients. J. Thromb. Thrombolysis 34(1), 120–125 (2012).
- 29. Yasumori T, Nagata K, Yang S *et al.* Cytochrome *P450* mediated metabolism of diazepam in human and rat: involvement of human CYP2C in N-demethylation in the substrate concentration-dependent manner. *Pharmacogenetics* 3(6), 291–301 (1993).
- 30. Brøsen K. Some aspects of genetic polymorphism in the biotransformation of antidepressants. Thérapie 59(1), 5-12 (2004).
- 31. Li X. Comparison of inhibitory effects of the proton pump-inhibiting drugs omeprazole, esomeprazole, lansoprazole, pantoprazole, and rabeprazole on human cytochrome P450 activities. *Drug Metab. Dispos.* 32(8), 821–827 (2004).
- 32. De Morais SM, Wilkinson GR, Blaisdell J, Nakamura K, Meyer UA, Goldstein JA. The major genetic defect responsible for the polymorphism of S-mephenytoin metabolism in humans. J. Biol. Chem. 269(22), 15419–15422 (1994).
- Hulot J. Cytochrome P450 2C19 loss-of-function polymorphism is a major determinant of clopidogrel responsiveness in healthy subjects. Blood 108(7), 2244–2247 (2006).
- 34. PharmVar CYP2C19. www.pharmvar.org/gene/CYP2C19
- 35. Sim S, Risinger C, Dahl M *et al.* A common novel *CYP2C19* gene variant causes ultrarapid drug metabolism relevant for the drug response to proton pump inhibitors and antidepressants. *Clin. Pharmacol. Ther.* 79(1), 103–113 (2006).
- 36. PharmGKB. CYP2C19 frequency table. https://api.pharmgkb.org/v1/download/file/attachment/CYP2C19_frequency_table.xlsx
- 37. Denisenko N, Sychev D, Sizova Z et al. Urine metabolic ratio of omeprazole in relation to CYP2C19 polymorphisms in Russian peptic ulcer patients. Pharmgenomics Pers. Med. 10, 253–259 (2017).
- Sychev DA, Denisenko NP, Sizova ZM, Grachev AV, Velikolug KA. The frequency of CYP2C19 genetic polymorphisms in Russian patients with peptic ulcers treated with proton pump inhibitors. *Pharmgenomics Pers. Med.* 8, 111–114 (2015).
- Mirzaev K, Zelenskaya E, Barbarash O et al. CYP2C19 polymorphism frequency in Russian patients in central Russia and Siberia with acute coronary syndrome. Pharmgenomics Pers. Med. 10, 107–114 (2017).
- 40. Golukhova E, Grigoryan M, Ryabinina M, Bulaeva N, Serebruany V. Body mass index and plasma P-selectin before coronary stenting predict high residual platelet reactivity at 6 months on dual antiplatelet therapy. *Cardiology* 139(2), 132–136 (2018).
- Zastrozhin M, Sorokin A, Agibalova T *et al.* Using a personalized clinical decision support system for bromdihydrochlorphenylbenzodiazepine dosing in patients with anxiety disorders based on the pharmacogenomic markers. *Hum. Psychopharmacol.* 33(6), e2677 (2018).
- 42. Bockeria O, Kudzoeva Z, Shvarts V, Koasari A, Donakanyan S. The possibility of selecting optimal antiplatelet therapy in patients with coronary heart disease in terms of *CYP2C19* polymorphism. *Terapevt. Arkh.* 88(5), 47–55 (2016).
- 43. Golukhova E, Ryabinina M, Bulaeva N, Grigorian M, Kubova M, Serebruany V. Clopidogrel response variability. *Am. J. Ther.* 22(3), 222–230 (2015).
- Khalikova A, Arkhipova A, Ahmetov I, Abdulkhakov R, Abdulkhakov S. [The study of cytochrome *P-450 CYP2C19* gene polymorphisms in population of Tatars living in Republic of Tatarstan]. *Pract. Med.* 3(58), 53–55 (2012) (In Russian).
- 45. Fedorinov D, Mirzaev K, Ivashchenko D *et al.* Pharmacogenetic testing by polymorphic markers 681G>A and 636G>A *CYP2C19* gene in patients with acute coronary syndrome and gastric ulcer in the Republic of Sakha (Yakutia). *Drug Metab. Pers. Ther.* 33(2), 91–98 (2018).
- 46. Muslimova E, Afanasiev S, Rebrova T, Sergienko T, Repin A. Association of *ITGB3*, *P2RY12*, and *CYP2C19* gene polymorphisms with platelet functional activity in patients with coronary heart disease during dual antiplatelet therapy. *Terapevt. Arkh.* 89(5), 74–85 (2017).
- 47. PharmVar CYP2D6. www.pharmvar.org/gene/CYP2D6

- 48. Owen R, Sangkuhl K, Klein T, Altman R. Cytochrome P450 2D6. Pharmacogenet. Genomics 19(7), 559-562 (2009).
- 49. Transon C, Leemann T, Dayer P. In vitro comparative inhibition profiles of major human drug metabolising cytochrome P450 isozymes (CYP2C9, CYP2D6 and CYP3A4) by HMG-CoA reductase inhibitors. Eur. J. Clin. Pharmacol. 50(3), 209-215 (1996).
- 50. Gardiner S. Pharmacogenetics, drug-metabolizing enzymes, and clinical practice. Pharmacol. Rev. 58(3), 521-590 (2006).
- 51. Ingelman-Sundberg M, Sim S, Gomez A, Rodriguez-Antona C. Influence of cytochrome P450 polymorphisms on drug therapies: pharmacogenetic, pharmacoepigenetic and clinical aspects. Pharmacol. Ther. 116(3), 496-526 (2007).
- 52. PharmGKB. CYP2D6 frequencies. https://api.pharmgkb.org/v1/download/file/attachment/CYP2D6_frequencies.xlsx
- 53. Sychev D, Zastrozhin M, Smirnov V, Grishina E, Savchenko L, Bryun E. The correlation between CYP2D6 isoenzyme activity and haloperidol efficacy and safety profile in patients with alcohol addiction during the exacerbation of the addiction. Pharmgenomics Pers. Med. 9, 89-95 (2016).
- 54. Zateyshchikov D, Minushkina L, Brovkin A et al. Association of CYP2D6 and ADRB1 genes with hypotensive and antichronotropic action of betaxolol in patients with arterial hypertension. Fundam Clin. Pharmacol. 21(4), 437-443 (2007).
- 55. Zastrozhin M, Grishina E, Denisenko N et al. Effects of CYP2D6 genetic polymorphisms on the efficacy and safety of fluvoxamine in patients with depressive disorder and comorbid alcohol use disorder. Pharmgenomics Pers. Med. 11, 113-119 (2018).
- 56. Sychev D, Zastrozhin M, Miroshnichenko I et al. Genotyping and phenotyping of CYP2D6 and CYP3A isoenzymes in patients with alcohol use disorder: correlation with haloperidol plasma concentration. Drug Metab. Pers. Ther. 32(3), 129-136 (2017).
- 57. Tiili E, Antikainen M, Mitiushkina N, Sukhovskaya O, Imyanitov E, Hirvonen A. Effect of genotype and methylation of CYP2D6 on smoking behaviour. Pharmacogenet. Genomics 25(11), 531-40 (2015).
- 58. Goryachkina K, Burbello A, Boldueva S, Babak S, Bergman U, Bertilsson L. CYP2D6 is a major determinant of metoprolol disposition and effects in hospitalized Russian patients treated for acute myocardial infarction. Eur. J. Clin. Pharmacol. 64(12), 1163-1173 (2008).
- 59. Goryachkina K, Burbello A, Boldueva S, Babak S, Bergman U, Bertilsson L. Inhibition of metoprolol metabolism and potentiation of its effects by paroxetine in routinely treated patients with acute myocardial infarction (AMI). Eur. J. Clin. Pharmacol. 64(3), 275-282 (2007)
- 60. Sychev D, Burashnikova I, Kazakov R. 1846G>A polymorphism of CYP2D6 gene and extrapyramidal side effects during antipsychotic therapy among Russians and Tatars: a pilot study. Drug Metab. Pers. Ther. 31(4), 205-212 (2016).
- 61. Polonikov A, Ivanov V, Solodilova M. Genetic variation of genes for xenobiotic-metabolizing enzymes and risk of bronchial asthma: the importance of gene-gene and gene-environment interactions for disease susceptibility. J. Hum. Genet. 54(8), 440-449 (2009).
- 62. Ivanova S, Filipenko M, Vyalova N et al. CYP1A2 and CYP2D6 gene polymorphisms in schizophrenic patients with neuroleptic drug-induced side effects. B. Exp. Biol. Med. 160(5), 687-690 (2016).
- 63. Fedorinov D, Mirzaev K, Mustafina V et al. Pharmacogenetic testing by polymorphic markers G1846A (CYP2D6*4) and C100T (CYP2D6*10) of the CYP2D6 gene in coronary heart disease patients taking ββ-blockers in the Republic of Sakha (YAKUTIA). Drug Metab. Pers. Ther. 33(4), 195-200 (2018).
- 64. Duzhak T, Mitrofanov D, Ostashevskii V et al. Genetic polymorphisms of CYP2D6, CYP1A1, GSTM1 and p53 genes in a unique Siberian population of Tundra Nentsi. Pharmacogenetics 10(6), 531-537 (2010).
- 65. PharmVar CYP1A1. www.pharmvar.org/gene/CYP1A1
- 66. Gurusamy U, Shewade D. Pharmacogenomics in India. Handbook Pharmacogenom. Strat. Med. Elsevier, PA, USA, 1037-1059 (2014).
- 67. Salnikova L, Smelaya T, Golubev A, Rubanovich A, Moroz V. CYP1A1, GCLC, AGT, AGTR1 gene-gene interactions in community-acquired pneumonia pulmonary complications. Mol. Biol. Rep. 40(11), 6163-6176 (2013).
- Salnikova L, Smelaya T, Moroz V, Golubev A, Rubanovich A. Functional polymorphisms in the CYPIA1, ACE, and IL-6 genes 68. contribute to susceptibility to community-acquired and nosocomial pneumonia. Int. J. Infect. Dis. 17(6), e433-e442 (2013).
- 69. Belogubova E, Ulibina Y, Suvorova I et al. Combined CYP1A1/GSTM1 at-risk genotypes are overrepresented in squamous cell lung carcinoma patients but underrepresented in elderly tumor-free subjects. J. Cancer Res. Clin. Oncol. 132(5), 327-331 (2006).
- 70. Mikhailova O, Gulyaeva L, Prudnikov A, Gerasimov A, Krasilnikov S. Estrogen-metabolizing gene polymorphisms in the assessment of female hormone-dependent cancer risk. Pharmacogenomics J. 6(3), 189-193 (2006).
- 71. PharmVar CYP2C8. www.pharmvar.org/gene/CYP2C8
- 72. Totah R, Rettie A. Cytochrome P450 2C8: substrates, inhibitors, pharmacogenetics, and clinical relevance. Clin. Pharmacol. Ther. 77(5), 341-352 (2005).
- 73. Bahadur N, Leathart J, Mutch E et al. CYP2C8 polymorphisms in Caucasians and their relationship with paclitaxel 6α-hydroxylase activity in human liver microsomes. Biochem. Pharmacol. 64(11), 1579-1589 (2002).
- Kurdyukov I, Shmurak V, Nadeyev A, Voitenko N, Prokofyeva D, Goncharov N. Esterase status of the organism at exposure to toxic 74. substances and pharmaceutical preparations. Toxicol. Rev. (6), 6-12 (2012).



- 75. Sychev D, Levanov A, Shelekhova T *et al.* The impact of *ABCB1* (rs1045642 and rs4148738) and *CES1* (rs2244613) gene polymorphisms on dabigatran equilibrium peak concentration in patients after total knee arthroplasty. *Pharmgenomics Pers. Med.* 11, 127–137 (2018).
- 76. Raftogianis R, Creveling C, Weinshilboum R, Weisz J. Estrogen metabolism by conjugation. J. Natl Cancer Inst. Monogr. 27, 113–124 (2000).
- 77. Sychev D, Rozhkov A, Kazakov R, Ananichuk A. The impact of *CYP4F2*, *ABCB1*, and *GGCX* polymorphisms on bleeding episodes associated with acenocoumarol in Russian patients with atrial fibrillation. *Drug Metab. Pers. Ther.* 31(3), 173–178 (2016).
- 78. Kryukov A, Sychev D, Andreev D *et al.* Influence of *ABCB1* and *CYP3A5* gene polymorphisms on pharmacokinetics of apixaban in patients with atrial fibrillation and acute stroke. *Pharmgenomics Pers. Med.* 11, 43–49 (2018).
- 79. Sychev D, Shikh N, Morozova T, Grishina E, Ryzhikova K, Malova E. Effects of *ABCB1* rs1045642 polymorphisms on the efficacy and safety of amlodipine therapy in Caucasian patients with stage I-II hypertension. *Pharmgenomics Pers. Med.* 11, 157–165 (2018).
- Shuev G, Sychev D, Chertovskih J, Grachev A, Syrkova O, Maksimova N. The frequency of SLCO1B1*5 polymorphism genotypes among Russian and Sakha (Yakutia) patients with hypercholesterolemia. *Pharmgenomics Pers. Med.* 25(9), 59–63 (2016).
- Rozhkov A, Sychev D, Kazakov R. ABCB1 polymorphism and acenocoumarol safety in patients with valvular atrial fibrillation. Int. J. Risk Saf. Med. 27(s1), S15–S16 (2015).
- Khokhlov A, Sychev D, Sirotkina A. [Safety aspects of using of statins: drug interaction, pharmacogenetic questions]. *Med. Pharmacol.* 1–2 (24) (2016) (in Russian).
- Solodun M, Yakushin S. Aspects of lipid-lowering therapy with atorvastatin in patients with myocardial infarction from the perspective of personalized medicine. *Ration Pharmacother. Cardiol.* 11(1), 31–35 (2015).
- Osswald E, Johne A, Laschinski G et al. Association of MDR1 genotypes with susceptibility to colorectal cancer in older non-smokers. Eur. J. Clin. Pharmacol. 63(1), 9–16 (2006).
- 85. Aszalos A. Drug-drug interactions affected by the transporter protein, P-glycoprotein (ABCB1, MDR1). Drug Discov. Today. 12(19–20), 838–843 (2007).
- Fung KL, Gottesman MM. A synonymous polymorphism in a common MDR1 (*ABCB1*) haplotype shapes protein function. *Biochim. Biophys. Acta* 1794, 860–871 (2009).
- Search Collaborative Group, Link E, Parish S et al. SLCO1B1 variants and statin-induced myopathy a genomewide study. N. Engl. J. Med. 359(8), 789–799 (2008).
- Becquemont L, Alfirevic A, Amstutz U *et al.* Practical recommendations for pharmacogenomics-based prescription: 2010 ESF–UB Conference on Pharmacogenetics and Pharmacogenomics. *Pharmacogenomics* 12(1), 113–124 (2011).
- 89. Luzum J, Cheung J. Does cardiology hold pharmacogenetics to an inconsistent standard? A comparison of evidence among recommendations. *Pharmacogenomics* 19(15), 1203–1216 (2018).
- 90. CDC. Working Group: methods/EGAPP/CDC. www.cdc.gov/egappreviews/methods.html.
- 91. Tuteja S, Limdi N. Pharmacogenetics in cardiovascular medicine. Curr. Genet. Med. Rep. 4(3), 119-129 (2016).
- 92. Haga S, Moaddeb J. Comparison of delivery strategies for pharmacogenetic testing services. *Pharmacogenet. Genomics* 24(3), 139–145 (2014).